



Multiple View Geometry: Solution Exercise Sheet 2

Prof. Dr. Daniel Cremers, Julia Bergbauer, Jakob Engel, TU Munich
<http://vision.in.tum.de/teaching/ss2014/mvg2014>

Part I: Theory

$$1. \lambda_a = \frac{(\lambda_a v_a)^T v_b}{\langle v_a, v_b \rangle} = \frac{v_a^T A^T v_b}{\langle v_a, v_b \rangle} = \frac{v_a^T A v_b}{\langle v_a, v_b \rangle} = \frac{v_a^T (\lambda_b v_b)}{\langle v_a, v_b \rangle} = \lambda_b$$

2. Let V be the orthonormal matrix (i.e. $V^T = V^{-1}$) given by the eigenvectors, and Σ the diagonal matrix containing the eigenvalues:

$$V = \begin{pmatrix} | & & | \\ v_1 & \cdots & v_n \\ | & & | \end{pmatrix} \quad \text{and} \quad \Sigma = \begin{pmatrix} \lambda_1 & 0 & \cdots \\ 0 & \ddots & 0 \\ \cdots & 0 & \lambda_n \end{pmatrix}.$$

As V is a basis, we can express x as a linear combination of the eigenvectors $x = V\alpha$ with $\alpha \in \mathbb{R}^n$, with $\sum_i \alpha_i^2 = \alpha^T \alpha = x^T V^T V x = x^T x = 1$. This gives

$$\begin{aligned} x^T A x &= x^T V \Sigma V^{-1} x \\ &= \alpha^T V^T V \Sigma V^T V \alpha \\ &= \alpha^T \Sigma \alpha = \sum_i \alpha_i^2 \lambda_i \end{aligned}$$

Considering $\sum_i \alpha_i^2 = 1$, we can conclude that this expression is minimized iff only the α_i corresponding to the smallest eigenvalue(s) are non-zero. If $\lambda_{n-1} \gtrsim \lambda_n$, there exist only two solutions ($\alpha_n = \pm 1$), otherwise infinitely many.

For maximisation, only the the α_i corresponding to the largest eigenvalue(s) can be non-zero.

3. We show that: $x \in \text{kernel}(A) \Leftrightarrow x \in \text{kernel}(A^T A)$.

$$\text{''}\Rightarrow\text{''}: \text{Let } x \in \text{kernel}(A) \\ A^T \underbrace{Ax}_{=0} = A^T 0 = 0 \Rightarrow x \in \text{kernel}(A^T A)$$

$$\text{''}\Leftarrow\text{''}: \text{Let } x \in \text{kernel}(A^T A) \\ 0 = x^T \underbrace{A^T Ax}_{=0} = \langle Ax, Ax \rangle = \|Ax\|^2 \Rightarrow Ax = 0 \Rightarrow x \in \text{kernel}(A)$$