

Chapter 1

Convex Analysis

Convex Optimization for Machine Learning & Computer Vision
SS 2017

Tao Wu
Thomas Möllenhoff
Emanuel Laude

Computer Vision Group
Department of Informatics
TU München

Convex Analysis

Tao Wu
Thomas Möllenhoff
Emanuel Laude



Convex Set

Convex Function

Existence of Minimizer

Subdifferential

Convex Conjugate



Convex Set



Assume

- \mathbb{E} is a Euclidean space (finite dimensional vector space), equipped with the inner product $\langle \cdot, \cdot \rangle$, e.g. $\langle u, v \rangle = u \cdot v$.
- C is a closed convex subset in \mathbb{E} .
- J is a convex objective function.

Convex Set

Convex Function

Existence of Minimizer

Subdifferential

Convex Conjugate

Convex optimization

$$\text{minimize } J(u) \quad \text{over } u \in C.$$

First questions:

- What is a convex set?
- What is a convex function?

Convex set

Definition

A set C is said to be **convex** if

$$\alpha u + (1 - \alpha)v \in C, \quad \forall u, v \in C, \quad \forall \alpha \in [0, 1].$$



Convex Set


Convex Function

Existence of Minimizer


Subdifferential

Convex Conjugate

convex



non-convex



Definition

- A set $C \subset \mathbb{E}$ is **open** if $\forall u \in C, \exists \epsilon > 0$ s.t. $B_\epsilon(u) \subset C$, where $B_\epsilon(u) := \{v \in \mathbb{E} : \|v - u\| < \epsilon\}$.
- A set $C \subset \mathbb{E}$ is **closed** if its complement $\mathbb{E} \setminus C$ is open.
- The **closure** of a set $C \subset \mathbb{E}$ is

$$\text{cl } C = \{u \in \mathbb{E} : \exists \{u^k\} \subset C \text{ s.t. } \lim_{k \rightarrow \infty} u^k = u\}.$$

- The **interior** of a set $C \subset \mathbb{E}$ is

$$\text{int } C = \{u \in C : \exists \epsilon > 0 \text{ s.t. } B_\epsilon(u) \subset C\}.$$



Convex Set

Convex Function

Existence of Minimizer

Subdifferential

Convex Conjugate

Definition

- A set $C \subset \mathbb{E}$ is **open** if $\forall u \in C, \exists \epsilon > 0$ s.t. $B_\epsilon(u) \subset C$, where $B_\epsilon(u) := \{v \in \mathbb{E} : \|v - u\| < \epsilon\}$.
- A set $C \subset \mathbb{E}$ is **closed** if its complement $\mathbb{E} \setminus C$ is open.
- The **closure** of a set $C \subset \mathbb{E}$ is

$$\text{cl } C = \{u \in \mathbb{E} : \exists \{u^k\} \subset C \text{ s.t. } \lim_{k \rightarrow \infty} u^k = u\}.$$

- The **interior** of a set $C \subset \mathbb{E}$ is

$$\text{int } C = \{u \in C : \exists \epsilon > 0 \text{ s.t. } B_\epsilon(u) \subset C\}.$$

- The **relative interior** of a convex set $C \subset \mathbb{E}$ is

$$\text{ri } C = \{u \in C : \forall v \in C, \exists \alpha > 1 \text{ s.t. } v + \alpha(u - v) \in C\}.$$



Convex Set

Convex Function

Existence of Minimizer

Subdifferential

Convex Conjugate



The following operations preserve the convexity:

- Intersection: $C_1 \cap C_2$
- Summation: $C_1 + C_2 := \{u^1 + u^2 : u^1 \in C_1, u^2 \in C_2\}$
- Closure: $\text{cl } C$
- Interior: $\text{int } C$

– The union of convex sets is not convex in general.



The following operations preserve the convexity:

- Intersection: $C_1 \cap C_2$
- Summation: $C_1 + C_2 := \{u^1 + u^2 : u^1 \in C_1, u^2 \in C_2\}$
- Closure: $\text{cl } C$
- Interior: $\text{int } C$


– The union of convex sets is not convex in general.

– *Polyhedral sets* are always convex; *cones* are not necessarily convex.

Convex cone

C is a **cone** if $C = \alpha C$ for any $\alpha > 0$. C is a **convex cone** if C is a cone and is convex as well.

Separation of convex sets



Theorem (separation of convex sets)

Let C_1, C_2 be nonempty convex subsets in \mathbb{E} s.t. $C_1 \cap C_2 = \emptyset$ and C_1 is open. Then there exists a hyperplane separating C_1 and C_2 , i.e. $\exists v \in \mathbb{E}, v \neq 0, \alpha \in \mathbb{R}$ s.t.

$$\langle v, u^1 \rangle \geq \alpha \geq \langle v, u^2 \rangle, \quad \forall u^1 \in C_1, u^2 \in C_2.$$

Proof: on board.





Theorem (separation of convex sets)

Let C_1, C_2 be nonempty convex subsets in \mathbb{E} s.t. $C_1 \cap C_2 = \emptyset$ and C_1 is open. Then there exists a hyperplane separating C_1 and C_2 , i.e. $\exists v \in \mathbb{E}, v \neq 0, \alpha \in \mathbb{R}$ s.t.

$$\langle v, u^1 \rangle \geq \alpha \geq \langle v, u^2 \rangle, \quad \forall u^1 \in C_1, u^2 \in C_2.$$

Proof: on board.

Remarks

- 1 The proof works in any real Hilbert space.
- 2 Corollary: In a real Hilbert space, any (strongly) closed convex subset C is weakly closed.

Convex Set

Convex Function

Existence of Minimizer

Subdifferential

Convex Conjugate



Convex Set

Convex Function

Existence of Minimizer

Subdifferential

Convex Conjugate

Convex Function



- An **extended real-valued function** J maps from \mathbb{E} to $\overline{\mathbb{R}} := \mathbb{R} \cup \{\infty\}$.
- The **domain** of $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ is

$$\text{dom } J = \{u \in \mathbb{E} : J(u) < \infty\}.$$

- The function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ is **proper** if $\text{dom } J \neq \emptyset$.

Definition

We say $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ is a **convex function** if

- 1 $\text{dom } J$ is a convex set.
- 2 For all $u, v \in \text{dom } J$ and $\alpha \in [0, 1]$ it holds that

$$J(\alpha u + (1 - \alpha)v) \leq \alpha J(u) + (1 - \alpha)J(v).$$

We say J is **strictly convex** if the above inequality is strict for all $\alpha \in (0, 1)$ and $u \neq v$.

Examples

- $J_{data}(u) = \|u - f\|_q^q$ where $q \geq 1$ and $\|\cdot\|_q$ is the ℓ^q -norm.
- $J_{regu}(u) = \|Ku\|_q^q$ where K is linear transform.
- $J(u) = J_{data}(u) + \alpha J_{regu}(u)$ where $\alpha > 0$.





- $J_{data}(u) = \|u - f\|_q^q$ where $q \geq 1$ and $\|\cdot\|_q$ is the ℓ^q -norm.
- $J_{regu}(u) = \|Ku\|_q^q$ where K is linear transform.
- $J(u) = J_{data}(u) + \alpha J_{regu}(u)$ where $\alpha > 0$.
- **Indicator function:**

$$\delta_C(u) = \begin{cases} 0 & \text{if } u \in C, \\ \infty & \text{otherwise,} \end{cases}$$

where C is a convex subset of \mathbb{E} .

- Alternative formulation of constrained optimization:

$$\min J(u) \text{ over } u \in C. \Leftrightarrow \min J(u) + \delta_C(u) \text{ over } u \in \mathbb{E}.$$



(As exercises)

- Any norm (over a normed vector space) is a convex function.
- J is a convex function and K is a linear transform $\Rightarrow J(K\cdot)$ is convex function.
- (Jensen's inequality) $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ is convex iff

$$J\left(\sum_{i=1}^n \alpha_i u^i\right) \leq \sum_{i=1}^n \alpha_i J(u^i),$$


whenever $\{u^i\}_{i=1}^n \subset \mathbb{E}$, $\{\alpha_i\}_{i=1}^n \subset [0, 1]$, $\sum_{i=1}^n \alpha_i = 1$.

Epigraph

Definition

The **epigraph** of a proper function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ is

$$\text{epi } J = \{(u, \alpha) \in \mathbb{E} \times \mathbb{R} : J(u) \leq \alpha\}.$$



Theorem

A proper function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ is convex (resp. strictly convex) iff $\text{epi } J$ is a convex (resp. strictly convex) set.

Proof: as exercise.





Definition

Assume $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ with $\text{ri dom } J \neq \emptyset$. We say J is **locally Lipschitz** at $u \in \text{ri dom } J$ with modulus $L_u > 0$ if there exists $\epsilon > 0$ s.t.

$$|J(u^1) - J(u^2)| \leq L_u \|u^1 - u^2\| \quad \forall u^1, u^2 \in B_\epsilon(u) \cap \text{ri dom } J.$$

[Convex Set](#)[Convex Function](#)[Existence of Minimizer](#)[Subdifferential](#)[Convex Conjugate](#)



Definition

Assume $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ with $\text{ri dom } J \neq \emptyset$. We say J is **locally Lipschitz** at $u \in \text{ri dom } J$ with modulus $L_u > 0$ if there exists $\epsilon > 0$ s.t.

$$|J(u^1) - J(u^2)| \leq L_u \|u^1 - u^2\| \quad \forall u^1, u^2 \in B_\epsilon(u) \cap \text{ri dom } J.$$

Theorem

A proper convex function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ is locally Lipschitz at any $u \in \text{ri dom } J$.

Proof: on board.

[Convex Set](#)[Convex Function](#)[Existence of Minimizer](#)[Subdifferential](#)[Convex Conjugate](#)

Global vs. Local minimizers

Recall the optimization of $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$:

$$\text{minimize } J(u) \quad \text{over } u \in \mathbb{E}.$$

Definition

- 1 $u^* \in \mathbb{E}$ is a **global minimizer** if $J(u^*) \leq J(u)$ for all $u \in \mathbb{E}$.
- 2 u^* is a **local minimizer** if $\exists \epsilon > 0$ s.t. $J(u^*) \leq J(u)$ for all $u \in B_\epsilon(u^*)$.
- 3 In the above definitions, a global/local minimizer is **strict** if $J(u^*) \leq J(u)$ is replaced by strict inequality $J(u^*) < J(u)$.



Global vs. Local minimizers

Recall the optimization of $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$:

$$\text{minimize } J(u) \quad \text{over } u \in \mathbb{E}.$$

Definition

- 1 $u^* \in \mathbb{E}$ is a **global minimizer** if $J(u^*) \leq J(u)$ for all $u \in \mathbb{E}$.
- 2 u^* is a **local minimizer** if $\exists \epsilon > 0$ s.t. $J(u^*) \leq J(u)$ for all $u \in B_\epsilon(u^*)$.
- 3 In the above definitions, a global/local minimizer is **strict** if $J(u^*) \leq J(u)$ is replaced by strict inequality $J(u^*) < J(u)$.

Theorem

For any proper convex function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$, if $u^* \in \text{dom } J$ is a local minimizer of J , then it is also a global minimizer.

Proof: on board.





Convex Set

Convex Function

Existence of Minimizer

Subdifferential

Convex Conjugate

Existence of Minimizer

Does a minimizer always exist?






- Consider

$$\text{minimize } J(u) \quad \text{over } u \in \mathbb{E},$$

where $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ is a proper, convex function.

- Some counterexamples for $J : \mathbb{R} \rightarrow \overline{\mathbb{R}}$:





- We shall formalize our observations and derive sufficient conditions for existence.

Sufficient conditions for existence

Definition

- 1 J is **bounded from below** if $J(\cdot) \geq C$ for some $C \in \mathbb{R}$.
- 2 J is **coercive** if $J(u) \rightarrow \infty$ whenever $\|u\| \rightarrow \infty$.
 - Proposition: J is coercive if $\text{dom } J$ is bounded.
- 3 J is **lower-semicontinuous** (l.s.c.) if

$$J(u^*) \leq \liminf_{u \rightarrow u^*} J(u).$$



- Proposition: J is l.s.c. iff $\text{epi } J$ is closed.



Sufficient conditions for existence

Definition

- 1 J is **bounded from below** if $J(\cdot) \geq C$ for some $C \in \mathbb{R}$.
- 2 J is **coercive** if $J(u) \rightarrow \infty$ whenever $\|u\| \rightarrow \infty$.
- 3 J is **lower-semicontinuous** (l.s.c.) if

$$J(u^*) \leq \liminf_{u \rightarrow u^*} J(u).$$

Theorem

Any proper function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$, which is bounded from below, coercive, and l.s.c., has a (global) minimizer.

Proof: on board.



Sufficient conditions for existence

Definition

- 1 J is **bounded from below** if $J(\cdot) \geq C$ for some $C \in \mathbb{R}$.
- 2 J is **coercive** if $J(u) \rightarrow \infty$ whenever $\|u\| \rightarrow \infty$.
- 3 J is **lower-semicontinuous** (l.s.c.) if

$$J(u^*) \leq \liminf_{u \rightarrow u^*} J(u).$$

Theorem

Any proper function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$, which is bounded from below, coercive, and l.s.c., has a (global) minimizer.

Proof: on board.

Remarks for infinite dimensions

- 1 Weak compactness in reflexive Banach (e.g. Hilbert) sp.
- 2 J is convex and strongly l.s.c. $\Rightarrow J$ is weakly l.s.c.





- Recall that a function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ is strictly convex if

$$J(\alpha u + (1 - \alpha)v) < \alpha J(u) + (1 - \alpha)J(v),$$

for all $u, v \in \text{dom } J$, $u \neq v$, $\alpha \in (0, 1)$.

Convex Set

Convex Function

Existence of Minimizer

Subdifferential

Convex Conjugate

Theorem

The minimizer of a strictly convex function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ is unique.

Proof: on board.



Subdifferential


Convex Set

Convex Function

Existence of Minimizer

Subdifferential

Convex Conjugate




Definition

$J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ is called (Fréchet) **differentiable** at $u \in \text{ri dom } J$ and $\nabla J(u) \in \mathbb{E}$ is the (Fréchet) **differential** of J at u if

$$\lim_{h \rightarrow 0, h \in (\text{dom } J) - u} \frac{J(u + h) - J(u) - \langle \nabla J(u), h \rangle}{\|h\|} = 0.$$

J is said **continuously differentiable** at $u \in \text{ri dom } J$ if $\nabla J(\cdot)$ is continuous on $\text{dom } J \cap B_\epsilon(u)$ for some $\epsilon > 0$.





Definition

$J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ is called (Fréchet) **differentiable** at $u \in \text{ri dom } J$ and $\nabla J(u) \in \mathbb{E}$ is the (Fréchet) **differential** of J at u if

$$\lim_{h \rightarrow 0, h \in (\text{dom } J) - u} \frac{J(u+h) - J(u) - \langle \nabla J(u), h \rangle}{\|h\|} = 0.$$

J is said **continuously differentiable** at $u \in \text{ri dom } J$ if $\nabla J(\cdot)$ is continuous on $\text{dom } J \cap B_\epsilon(u)$ for some $\epsilon > 0$.

Remark

In functional analysis, $\nabla J(u)$ is treated as a *dual* object in \mathbb{E}^* , and $\langle \nabla J(u), h \rangle_{\mathbb{E}^*, \mathbb{E}}$ as *duality pairing*.



Convex Set

Convex Function


Existence of Minimizer

Subdifferential

Convex Conjugate

Subdifferential

Now we generalize differentiability from differentiable functions to nonsmooth (convex) functions.



Definition


The **subdifferential** of a convex function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ at $u \in \text{dom } J$ is defined by

$$\partial J(u) = \{ \xi \in \mathbb{E} : J(v) \geq J(u) + \langle \xi, v - u \rangle \}.$$



Subdifferential

Now we generalize differentiability from differentiable functions to nonsmooth (convex) functions.



Definition

The **subdifferential** of a convex function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ at $u \in \text{dom } J$ is defined by

$$\partial J(u) = \{ \xi \in \mathbb{E} : J(v) \geq J(u) + \langle \xi, v - u \rangle \}.$$

Geometric interpretation

$\xi \in \partial J(u)$ iff $(\xi, -1)$ is a normal vector for the supporting hyperplane of $\text{epi } J$ at $(u, J(u))$.



Basic facts

- 1 $\partial J(\cdot)$ is a **set-valued map**.
- 2 If J is cont. differentiable at u , then $\partial J(u) = \{\nabla J(u)\}$.



Convex Set

Convex Function

Existence of Minimizer

Subdifferential

Convex Conjugate



Basic facts

- 1 $\partial J(\cdot)$ is a **set-valued map**.
- 2 If J is cont. differentiable at u , then $\partial J(u) = \{\nabla J(u)\}$.

Examples (as exercises)

- 1 $u \in \mathbb{R}^n \mapsto \|u\|_1$.
- 2 $u \in \mathbb{R}^n \mapsto \|u\|_\infty$.
- 3 $X \in \mathbb{R}^{n \times n} \mapsto \|X\|_{1,2} = \sum_i \left(\sum_j |X_{i,j}|^2 \right)^{1/2}$.
- 4 $X \in \mathbb{R}^{n \times n} \mapsto \|X\|_{nuc} = \sum_i \sigma_i(X)$ (sum of singular values).
- 5 Given a closed convex subset $C \subset \mathbb{E}$ and $u \in C$,

$$\partial \delta_C(u) = N_C(u) = \{\xi \in \mathbb{E} : \langle \xi, v - u \rangle \leq 0 \forall v \in C\},$$

known as the **normal cone** of C at u .

Convex Set

Convex Function

Existence of Minimizer

Subdifferential

Convex Conjugate

Theorem (chain rule under linear transform)

Let $\tilde{J}(\cdot) = J(K\cdot)$ with some convex function J and linear transform K . Then

$$\partial\tilde{J}(u) = K^T \partial J(Ku)$$

for any $u \in \text{ri dom } J$.

Example: $J(u) = \|Ku\|_1 \Rightarrow \partial J(u) = K^T \partial \|\cdot\|_1(Ku)$.



Convex Set

Convex Function

Existence of Minimizer

Subdifferential

Convex Conjugate

Theorem (chain rule under linear transform)

Let $\tilde{J}(\cdot) = J(K\cdot)$ with some convex function J and linear transform K . Then

$$\partial\tilde{J}(u) = K^\top \partial J(Ku)$$

for any $u \in \text{ri dom } J$.

Example: $J(u) = \|Ku\|_1 \Rightarrow \partial J(u) = K^\top \partial \|\cdot\|_1(Ku)$.

Theorem (summation rule)

If $J(\cdot) = J_1(\cdot) + J_2(\cdot)$ for some convex functions J_1 and J_2 , then

$$\partial\tilde{J}(u) = \partial J_1(u) + \partial J_2(u)$$

for any $u \in \text{ri dom } J_1 \cap \text{ri dom } J_2$.

Warning: not true if J_1 or J_2 is non-convex, e.g. $0 = |\cdot| + (-|\cdot|)$.



Convex Set

Convex Function

Existence of Minimizer

Subdifferential

Convex Conjugate

Properties of subdifferential map

Theorem

Let $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ be a convex function. Then ∂J is a **monotone operator**, i.e. $\forall u^1, u^2 \in \text{dom } J, \xi^1 \in \partial J(u^1), \xi^2 \in \partial J(u^2)$:

$$\langle \xi^1 - \xi^2, u^1 - u^2 \rangle \geq 0.$$

Proof: on board.



Properties of subdifferential map

Theorem

Let $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ be a convex function. Then ∂J is a **monotone operator**, i.e. $\forall u^1, u^2 \in \text{dom } J, \xi^1 \in \partial J(u^1), \xi^2 \in \partial J(u^2)$:

$$\langle \xi^1 - \xi^2, u^1 - u^2 \rangle \geq 0.$$

Proof: on board.

Theorem

Let $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ be a convex function. Then for any $u \in \text{ri dom } J$, $\partial J(u)$ is a nonempty, compact, and convex subset.

Proof: on board.



Properties of subdifferential map

Theorem

Let $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ be a convex function. Then ∂J is a **monotone operator**, i.e. $\forall u^1, u^2 \in \text{dom } J, \xi^1 \in \partial J(u^1), \xi^2 \in \partial J(u^2)$:

$$\langle \xi^1 - \xi^2, u^1 - u^2 \rangle \geq 0.$$

Proof: on board.

Theorem

Let $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ be a convex function. Then for any $u \in \text{ri dom } J$, $\partial J(u)$ is a nonempty, compact, and convex subset.

Proof: on board.

Theorem

Let $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ be a proper, convex, l.s.c. function. Then the set-valued map $\partial J(\cdot)$ is **closed**, i.e. $\xi^* \in \partial J(u^*)$ whenever

$$\exists (u^k, \xi^k) \rightarrow (u^*, \xi^*) \in (\text{ri dom } J) \times \mathbb{E} \text{ s.t. } \xi^k \in \partial J(u^k) \forall k.$$

Proof: on board.



Optimality condition

Theorem

Given any proper convex function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$, the sufficient and necessary condition for u^* being a (global) minimizer for J is

$$0 \in \partial J(u^*).$$

Proof: on board.



Optimality condition

Theorem

Given any proper convex function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$, the sufficient and necessary condition for u^* being a (global) minimizer for J is


$$0 \in \partial J(u^*).$$

Proof: on board.

Constrained optimization as special case

If $\tilde{J}(u) = J(u) + \delta_C(u)$ with convex function $J : \mathbb{E} \rightarrow \mathbb{R}$ and closed convex subset $C \in \mathbb{E}$, then $0 \in \partial J(u^*) \Leftrightarrow$

$$0 \in \partial J(u^*) + N_C(u^*).$$



Optimality condition

Theorem

Given any proper convex function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$, the sufficient and necessary condition for u^* being a (global) minimizer for J is

$$0 \in \partial J(u^*).$$

Proof: on board.

Constrained optimization as special case

If $\tilde{J}(u) = J(u) + \delta_C(u)$ with convex function $J : \mathbb{E} \rightarrow \mathbb{R}$ and closed convex subset $C \in \mathbb{E}$, then $0 \in \partial J(u^*) \Leftrightarrow$

$$0 \in \partial J(u^*) + N_C(u^*).$$

Remark

$0 \in \partial J(u^*) + N_C(u^*)$ is a *geometric* optimality condition. Further characterization relies on the algebraic representation of $N_C(u^*)$, e.g. the Karush-Kuhn-Tucker (KKT) conditions under certain constraint qualifications.





Convex Conjugate

Convex Set

Convex Function

Existence of Minimizer

Subdifferential

Convex Conjugate

Legendre-Fenchel transform

Definition

Given a convex function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$, the **convex conjugate** of J is defined by

$$J^*(p) = \sup_{u \in \mathbb{E}} \langle u, p \rangle - J(u).$$



Legendre-Fenchel transform

Definition

Given a convex function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$, the **convex conjugate** of J is defined by

$$J^*(p) = \sup_{u \in \mathbb{E}} \langle u, p \rangle - J(u).$$

Examples

- 1 $J(\cdot) = \langle w, \cdot \rangle \Rightarrow J^*(\cdot) = \delta\{\cdot = w\}. (w \in \mathbb{E})$
- 2 $J(\cdot) = \frac{1}{q} \|\cdot\|_q^q \Rightarrow J^*(\cdot) = \frac{1}{q'} \|\cdot\|_{q'}^{q'}. (0 < q < 1, \frac{1}{q} + \frac{1}{q'} = 1)$
- 3 $J(\cdot) = \|\cdot\| \Rightarrow J^*(\cdot) = \delta\{\|\cdot\|_* \leq 1\}. (\|\cdot\|_* \text{ is the dual norm of } \|\cdot\|, \text{ i.e. } \|v\|_* = \sup_{\|u\| \leq 1} \langle u, v \rangle)$
- 4 $J(\cdot) = \delta\{\|\cdot\|_* \leq 1\} \Rightarrow J^*(\cdot) = \|\cdot\|.$



Legendre-Fenchel transform

Definition

Given a convex function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$, the **convex conjugate** of J is defined by

$$J^*(p) = \sup_{u \in \mathbb{E}} \langle u, p \rangle - J(u).$$

Examples

- 1 $J(\cdot) = \langle w, \cdot \rangle \Rightarrow J^*(\cdot) = \delta\{\cdot = w\}. (w \in \mathbb{E})$
- 2 $J(\cdot) = \frac{1}{q} \|\cdot\|_q^q \Rightarrow J^*(\cdot) = \frac{1}{q'} \|\cdot\|_{q'}^{q'}. (0 < q < 1, \frac{1}{q} + \frac{1}{q'} = 1)$
- 3 $J(\cdot) = \|\cdot\| \Rightarrow J^*(\cdot) = \delta\{\|\cdot\|_* \leq 1\}. (\|\cdot\|_* \text{ is the dual norm of } \|\cdot\|, \text{ i.e. } \|v\|_* = \sup_{\|u\| \leq 1} \langle u, v \rangle)$
- 4 $J(\cdot) = \delta\{\|\cdot\|_* \leq 1\} \Rightarrow J^*(\cdot) = \|\cdot\|.$


Basic facts

- Scalar multiplication: $\tilde{J}(\cdot) = \alpha J(\cdot) \Rightarrow \tilde{J}^*(\cdot) = \alpha J^*(\cdot/\alpha).$
- Translation: $\tilde{J}(\cdot) = J(\cdot - z) \Rightarrow \tilde{J}^*(\cdot) = J^*(\cdot) + \langle \cdot, z \rangle.$



Geometric interpretation

Convex conjugation maps from:
non-vertical hyperplanes supporting the epigraph
to:
crossing points of vertical axis.



Courtesy of Bertsekas





Theorem (Fenchel-Young inequality)

For all $u \in \text{dom } J$, $p \in \text{dom } J^*$, we have

$$J(u) + J^*(p) \geq \langle u, p \rangle.$$

The equality holds iff $p \in \partial J(u)$.

Proof: $J(u) + J^*(p) \geq \langle u, p \rangle$ follows directly from the definition of convex conjugate; $p \in \partial J(u)$ is the sufficient and necessary condition for: $u = \arg \min_{v \in \mathbb{E}} J(v) - \langle v, p \rangle$.

Convex Set

Convex Function

Existence of Minimizer

Subdifferential

Convex Conjugate



Theorem

Assume $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ and $J^{**} = (J^*)^*$ is the **biconjugate** of J .

In general:

- 1 $J^{**}(\cdot) \leq J(\cdot)$.
- 2 J^* is convex and l.s.c.

If J is proper, convex, and l.s.c., then:

- 3 $J^{**}(\cdot) = J(\cdot)$.
- 4 $p \in \partial J(u)$ iff $u \in \partial J^*(p)$.

Proof: on board.

[Convex Set](#)[Convex Function](#)[Existence of Minimizer](#)[Subdifferential](#)[Convex Conjugate](#)