

# Analysis of 3D Shapes (IN2238)



Frank R. Schmidt  
Matthias Vestner

Summer Semester 2017

## 2. LA and Analysis (Recap 1)

## Linear Spaces

## Shapes and Objects



The concept of a shape can be understood as a generalization of objects. In fact, we defined the **shape of an object** as a class of equivalent objects.

As a 3D-object we understand something like a ball or a human that occupies a certain region  $X \subset \mathbb{R}^3$  of the real world.

## Linear Algebra and Analysis



We defined an object of dimension  $d$  as an open subset  $X \subset \mathbb{R}^d$  such that  $\dim X = d - 1$ . This concept of a dimension is in fact an extension of the dimension for linear vector spaces.

As a matter of fact, we like this boundary to be smooth.

Therefore, we will recap some of the main concepts from **Linear Algebra** as well as **Analysis**.

## Vector Spaces

An  $\mathbb{R}$ -vector space  $V$  is formally defined as an **Abelian group** that has some additional **linear** properties.

As an Abelian group,  $V$  possesses a binary operation  $+$ , a neutral element  $0 \in V$  as well as an inverse element  $(-v) \in V$  for each element  $v \in V$ .

In addition, there exists a scalar multiplication  $\cdot$  such that

$$\begin{aligned} (\lambda \cdot \mu) \cdot v &= \lambda \cdot (\mu \cdot v) & \forall \lambda, \mu \in \mathbb{R}, v \in V \\ 1 \cdot v &= v & \forall v \in V \\ (\lambda + \mu) \cdot v &= \lambda \cdot v + \mu \cdot v & \forall \lambda, \mu \in \mathbb{R}, v \in V \\ \lambda \cdot (u + v) &= \lambda \cdot u + \lambda \cdot v & \forall \lambda \in \mathbb{R}, u, v \in V \end{aligned}$$

Examples for vector spaces are  $\mathbb{R}^n$ , but also **function spaces** like  $C^k(\mathbb{R}^n)$ .

## Vector Sub-Spaces

Given an  $\mathbb{R}$ -vector space  $V$ , the subset  $U \subset V$  is also a vector space (called **subspace**) if the following holds:

$$\begin{aligned} u + v &\in U & \forall u, v \in U \\ \lambda v &\in U & \forall \lambda \in \mathbb{R}, v \in U \end{aligned}$$

Given a subset  $X = \{x_1, \dots, x_n\} \subset V$ , the subset  $\text{span}(X)$  is a subspace with

$$\text{span}(X) := \left\{ \sum_{i=1}^n \lambda_i x_i \mid \lambda \in \mathbb{R}^n \right\}.$$

If the  $x_i$  are linear independent, we call  $X$  a **base** of  $\text{span}(X)$  and  $n$  is called its **dimension**.

## Linear Mapping

# Linear Mapping

Given the  $\mathbb{R}$ -vector spaces  $U$  and  $V$ , a mapping  $L: U \rightarrow V$  is a **linear mapping** if the following holds:

$$\begin{aligned} L(u+v) &= L(u) + L(v) & \forall u, v \in U \\ L(\lambda u) &= \lambda L(u) & \forall \lambda \in \mathbb{R}, u \in U \end{aligned}$$

Is  $X$  a basis of the  $n$ -dimensional vector space  $U$  and  $Y$  a basis of the  $m$ -dimensional vector space  $V$ , we obtain

$$L(x_j) = \sum_{i=1}^m a_{ij} y_i$$

$A \in \mathbb{R}^{m \times n}$  is then called the **representing matrix** of  $L$  with respect to the bases  $X$  and  $Y$  and we write:

$$\mathcal{M}_Y^X(L) = A.$$

# Matrix-Multiplication

Given matrices  $A \in \mathbb{R}^{m \times r}$  and  $B \in \mathbb{R}^{r \times n}$ , the product  $C := A \cdot B \in \mathbb{R}^{m \times n}$  is

$$\begin{pmatrix} a_{11} & \cdots & a_{1r} \\ \vdots & & \vdots \\ a_{i1} & \cdots & a_{ir} \\ \vdots & & \vdots \\ a_{m1} & \cdots & a_{mr} \end{pmatrix} \cdot \begin{pmatrix} b_{11} & \cdots & b_{1j} & \cdots & b_{1n} \\ \vdots & & \vdots & & \vdots \\ b_{r1} & \cdots & b_{rj} & \cdots & b_{rn} \end{pmatrix} = \begin{pmatrix} c_{11} & \cdots & c_{1n} \\ \vdots & & \vdots \\ \vdots & & c_{ij} \\ \vdots & & \vdots \\ c_{m1} & \cdots & c_{mn} \end{pmatrix}$$

with

$$c_{ij} = \sum_{k=1}^r a_{ik} b_{kj}$$

It turns out that

$$\mathcal{M}_Z^Y(L_2) \cdot \mathcal{M}_Y^X(L_1) = \mathcal{M}_Z^X(L_2 \circ L_1)$$

# Change of Bases

If we have  $V = \mathbb{R}^n$  and a matrix  $A \in \mathbb{R}^{n \times n}$ , the matrix-vector multiplication defines a linear mapping:

$$\begin{aligned} L: \mathbb{R}^n &\rightarrow \mathbb{R}^n \\ x &\mapsto Ax \end{aligned}$$

Let us assume that we want to **change the bases** of  $\mathbb{R}^n$ . To that end, both  $X$  and  $Y$  can be written in matrix form and we have

$$\mathcal{M}_Y^X(L) = Y \cdot A \cdot X^{-1}$$

Thus, there is a subtle difference between linear mappings  $L$  and matrices  $A$ .  $A$  is a representation of  $L$  that also takes the specific bases into account.

We say that two matrices  $A$  and  $B$  are **similar**, if there exists an invertible matrix  $X$  such that  $B = X \cdot A \cdot X^{-1}$ .

# Differential

# History of Differential


While the concept of the **derivative** or **differential** is nowadays one of the basic concepts in modern mathematics, it took a while to find a clean mathematical definition.

The notation  $\frac{dy}{dx}$  is due to **Leibniz** who called **dx** and **dy** an "infinitely small change of"  $x$  resp.  $y$ .

In 1924, **Courant** mentioned that the idea of the differential as infinite small expression "lacks any meaning" and is therefore "useless".

The modern notion of derivatives and differential is due to **Cauchy** and **Weierstraß**, which we want to revise in the following.

# Derivative according to Cauchy




The derivative  $f'(x_0)$  of a function  $f: \mathbb{R} \rightarrow \mathbb{R}$  at the position  $x_0 \in \mathbb{R}$  is

$$f'(x_0) := \lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

While this is a working mathematical definition, it is a bit difficult to extend it to arbitrary functions  $f: \mathbb{R}^n \rightarrow \mathbb{R}^m$ , since we cannot "divide by vectors".

# Differential according to Weierstraß



Given a function  $f: \mathbb{R} \rightarrow \mathbb{R}$  and a position  $x_0 \in \mathbb{R}$ , its **differential**  $Df(x_0)$  is the **unique linear mapping**  $L: \mathbb{R} \rightarrow \mathbb{R}$  such that

$$\begin{aligned} f(x_0 + h) &= f(x_0) + L[h] + r(h) \\ \lim_{h \rightarrow 0} \frac{r(h)}{|h|} &= 0 \end{aligned}$$

# Jacobi Matrix

Let  $f: \mathbb{R}^m \rightarrow \mathbb{R}^n$  be a differentiable function and  $x_0 \in \mathbb{R}^m$ . The differential

$$Df(x_0): \mathbb{R}^m \rightarrow \mathbb{R}^n$$

is a linear mapping.

Using the canonical bases  $\{e_1, \dots, e_m\}$  for  $\mathbb{R}^m$  and  $\{e_1, \dots, e_n\}$  for  $\mathbb{R}^n$ ,  $Df(x_0)$  can be written in matrix form, the **Jacobi matrix**

$$Df(x_0)[h] = J \cdot h \quad J = \begin{pmatrix} J_{1,1} & \cdots & J_{1,m} \\ \vdots & & \vdots \\ J_{n,1} & \cdots & J_{n,m} \end{pmatrix}$$

with

$$J_{i,j} = \langle e_i, J \cdot e_j \rangle = \langle e_i, Df(x_0)[e_j] \rangle = \lim_{h \rightarrow 0} \frac{f^i(x_0 + h \cdot e_j) - f^i(x_0)}{h} = \partial_j f^i(x_0)$$

## Chain Rule

Linear Spaces Linear Mapping Differential Two Theorems

Let  $f: \mathbb{R}^m \rightarrow \mathbb{R}^n$  and  $g: \mathbb{R}^k \rightarrow \mathbb{R}^m$  be differentiable functions. Then we have

$$\begin{aligned} (f \circ g)(x_0 + h) &= f(g(x_0) + Dg(x_0)[h] + r_g(h)) \\ &= (f \circ g)(x_0) + Df(g(x_0)) [Dg(x_0)[h] + r_g(h)] + \\ &\quad r_f(Dg(x_0)[h] + r_g(h)) \\ &= (f \circ g)(x_0) + Df(g(x_0)) [Dg(x_0)[h]] + r(h) \end{aligned}$$

Thus we have

$$D(\mathbf{f} \circ \mathbf{g})(x_0) = \mathbf{Df}(g(x_0)) \cdot \mathbf{Dg}(x_0)$$

IN2238 - Analysis of Three-Dimensional Shapes

2. Linear Algebra and Analysis (Recap 1) - 17 / 24

## Chain Rule (Example)

Linear Spaces Linear Mapping Differential Two Theorems

Let  $g_1, g_2: \mathbb{R}^m \rightarrow \mathbb{R}^n$  and  $f: \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  with  $f(x, y) := x + y$ , we have

$$\begin{aligned} D(g_1 + g_2)(x_0) &= D(f \circ g)(x_0) = Df(g(x_0)) \cdot Dg(x_0) \\ &= (\text{Id} \quad \text{Id}) \cdot \begin{pmatrix} Dg_1(x_0) \\ Dg_2(x_0) \end{pmatrix} = Dg_1(x_0) + Dg_2(x_0) \end{aligned}$$

Let  $g_1, g_2: \mathbb{R}^m \rightarrow \mathbb{R}$  and  $f: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$  with  $f(x, y) := x \cdot y$ , we have

$$\begin{aligned} D(g_1 \cdot g_2)(x_0) &= D(f \circ g)(x_0) = Df(g(x_0)) \cdot Dg(x_0) \\ &= (g_2(x_0) \quad g_1(x_0)) \cdot \begin{pmatrix} Dg_1(x_0) \\ Dg_2(x_0) \end{pmatrix} \\ &= Dg_1(x_0) \cdot g_2(x_0) + Dg_2(x_0) \cdot g_1(x_0) \end{aligned}$$

IN2238 - Analysis of Three-Dimensional Shapes

2. Linear Algebra and Analysis (Recap 1) - 18 / 24

## Two Theorems

Linear Spaces Linear Mapping Differential Two Theorems

If  $f, f^{-1}: \mathbb{R}^n \rightarrow \mathbb{R}^n$  are both continuously differentiable, we have

$$\begin{aligned} \text{Id} &= D[f \circ f^{-1}](x) = Df(f^{-1}(x)) \cdot D[f^{-1}](x) \\ D[f^{-1}](x) &= Df(f^{-1}(x))^{-1} \end{aligned}$$

Interestingly, also the opposite is (locally) true

**Theorem 1.** Let  $f: \mathbb{R}^n \rightarrow \mathbb{R}^n$  cont. differentiable and  $Df(x_0)$  invertible. Then there exist neighborhoods  $U(x_0)$  and  $V(y_0)$  with  $y_0 = f(x_0)$  such that

- $f: U \rightarrow V$  is a bijection.
- $f$  is continuously differentiable.
- $f^{-1}$  is continuously differentiable.

IN2238 - Analysis of Three-Dimensional Shapes

2. Linear Algebra and Analysis (Recap 1) - 20 / 24

## Inverse Functions as Implicit Functions

Linear Spaces Linear Mapping Differential Two Theorems

Usually, one defines the **square root function** in an implicit manner:

$$x - \sqrt{x^2} = 0$$

This can be formally done in the following way:

**Theorem 2 (Square Root).** Let  $\Phi: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$  be the function

$$\Phi(x, y) = x - y^2,$$

which satisfies  $\Phi(x_0, y_0) = 0$  for  $(x_0, y_0) = (4, 2)$ .

Then there exist neighborhoods  $U(4)$ ,  $V(2)$  and unique  $f: U \rightarrow V$  such that

- $f$  is continuously differentiable and  $f(4) = 2$ .
- $\Phi(x, f(x)) = 0$  for all  $x \in U$ .
- $f'(x) = -\frac{\partial_x \Phi(x, f(x))}{\partial_y \Phi(x, f(x))} = \frac{1}{2f(x)}$

IN2238 - Analysis of Three-Dimensional Shapes

2. Linear Algebra and Analysis (Recap 1) - 21 / 24

## Implicit Function Theorem

Linear Spaces Linear Mapping Differential Two Theorems

This can be generalized to the

**Theorem 3 (Implicit Function Theorem).** Let  $\Phi: \mathbb{R}^m \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  be a cont. differentiable mapping which satisfies  $\Phi(x_0, y_0) = 0$  for a  $(x_0, y_0) \in \mathbb{R}^{m+n}$  and  $\partial_y \Phi(x_0, y_0)$  is invertible.

Then there exist neighborhoods  $U(x_0)$ ,  $V(y_0)$  and a continuously differentiable function  $f: U \rightarrow V$  such that

$$\Phi(x, f(x)) = 0 \quad \forall x \in U$$

and


$$\begin{aligned} f(x_0) &= y_0 \\ f'(x) &= -(\partial_y \Phi(x, f(x)))^{-1} \partial_x \Phi(x, f(x)) \quad \forall x \in U \end{aligned}$$

IN2238 - Analysis of Three-Dimensional Shapes

2. Linear Algebra and Analysis (Recap 1) - 22 / 24

## Implicit Function Theorem (Example)

Linear Spaces Linear Mapping Differential Two Theorems



Note that we can also use the implicit function theorem if we are not looking for the inverse of a function.

The points  $p = (p_1, p_2, p_3)$  on the unit sphere satisfy

$$p_1^2 + p_2^2 + p_3^2 = 1$$

If we use the notation  $x = (p_1, p_2)$  and  $y = p_3$ , the requirements for the implicit function theorem are satisfied for  $x_0 = (0, 0)$ ,  $y_0 = 1$  as well as



$$\Phi(x, y) = \|x\|^2 + y^2 - 1$$

IN2238 - Analysis of Three-Dimensional Shapes

2. Linear Algebra and Analysis (Recap 1) - 23 / 24

## Implicit Function Theorem (Example)

Linear Spaces Linear Mapping Differential Two Theorems



In particular, there exists a neighborhood  $U(x_0) \subset \mathbb{R}^2$  and a mapping  $f: U \rightarrow \mathbb{R}$  such that

$$\begin{aligned} \varphi: U &\rightarrow \mathbb{R}^3 \\ x &\mapsto (x, f(x)) \end{aligned}$$

maps the 2D region  $U$  onto a part of the sphere.

Functions like  $\varphi$  can be used to map a subset of a two-dimensional linear space onto a subset of a two-dimensional curved space. These curved spaces are called **manifolds**.

IN2238 - Analysis of Three-Dimensional Shapes

2. Linear Algebra and Analysis (Recap 1) - 24 / 24

IN2238 - Analysis of Three-Dimensional Shapes

2. Linear Algebra and Analysis (Recap 1) - 24 / 24