


Analysis of 3D Shapes (IN2238)

Frank R. Schmidt
Matthias Vestner

Summer Semester 2017

4. Differential and Curvature	2
Differential	3
Derivative according to Cauchy	4
Differential according to Weierstraß	5
Jacobi Matrix	6
Chain Rule	7
Interpretation of the Differential	8
Push-Forward	9
Curve Representation of Tangent Vectors	10
Alternative Definition of Tangent Vectors	11
Differential as Push-Forward	12
Push-Forward of Submanifolds	13
Coordinate Interpretation	14
How to Compute the Differential	15
What is a Matrix? (Recap)	16
Matrix of the Differential	17

Curvature of 2D Objects	18
Planar Curves and Normals	19
Differential of the Normal Mapping	20
Derivative of the Normal Field	21
Curvature	22
Curvature of Implicit Submanifolds	23
Literature	24


Derivative according to Cauchy

The derivative $f'(x_0)$ of a function $f: \mathbb{R} \rightarrow \mathbb{R}$ at the position $x_0 \in \mathbb{R}$ is

$$f'(x_0) := \lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

While this is a working mathematical definition, it is a bit difficult to extend it to arbitrary functions $f: \mathbb{R}^n \rightarrow \mathbb{R}^m$, since we cannot “divide by vectors”.

Differential according to Weierstraß



Given a function $f: \mathbb{R} \rightarrow \mathbb{R}$ and a position $x_0 \in \mathbb{R}$, its differential $Df(x_0)$ is the **unique linear mapping** $L: \mathbb{R} \rightarrow \mathbb{R}$ such that

$$f(x_0 + h) = f(x_0) + L[h] + r(h)$$

$$\lim_{h \rightarrow 0} \frac{r(h)}{|h|} = 0$$

Jacobi Matrix

Let $f: \mathbb{R}^m \rightarrow \mathbb{R}^n$ be a differentiable function and $x_0 \in \mathbb{R}^m$. The differential

$$Df(x_0): \mathbb{R}^m \rightarrow \mathbb{R}^n$$

is a linear mapping.

Using the canonical bases $\{e_1, \dots, e_m\}$ for \mathbb{R}^m and $\{e_1, \dots, e_n\}$ for \mathbb{R}^n , $Df(x_0)$ can be written in matrix form, the **Jacobi matrix**

$$Df(x_0)[h] = J \cdot h \qquad J = \begin{pmatrix} J_{1,1} & \cdots & J_{1,m} \\ \vdots & & \vdots \\ J_{n,1} & \cdots & J_{n,m} \end{pmatrix}$$

with

$$J_{i,j} = \langle e_i, J \cdot e_j \rangle = \langle e_i, Df(x_0)[e_j] \rangle = \lim_{h \rightarrow 0} \frac{f^i(x_0 + h \cdot e_j) - f^i(x_0)}{h} = \partial_j f^i(x_0)$$

Chain Rule

Let $f: \mathbb{R}^m \rightarrow \mathbb{R}^n$ and $g: \mathbb{R}^k \rightarrow \mathbb{R}^m$ be differentiable functions. Then we have

$$\begin{aligned} (f \circ g)(x_0 + h) &= f(g(x_0) + Dg(x_0)[h] + r_g(h)) \\ &= (f \circ g)(x_0) + Df(g(x_0)) [Dg(x_0)[h] + r_g(h)] + \\ &\quad r_f(Dg(x_0)[h] + r_g(h)) \\ &= (f \circ g)(x_0) + Df(g(x_0)) [Dg(x_0)[h]] + r(h) \end{aligned}$$

Thus we have

$$D(\mathbf{f} \circ \mathbf{g})(x_0) = \mathbf{Df}(g(x_0)) \circ \mathbf{Dg}(x_0)$$

Interpretation of the Differential

Given a function $f: \mathbb{R}^m \rightarrow \mathbb{R}^n$ and a position $p \in \mathbb{R}^m$, the equation

$$f(p + v) = f(p) + Df(p)[v] + r(v)$$

can be interpreted as following:

- p describes a **point** in the space on which f is defined,
- v describes the **direction** in which we change the point p
- $Df(p)[v]$ describes the **direction** in which f changes if we change the point p in the direction v .

For vector spaces, there is no distinction between points and directions. For manifolds M , points p will be on the manifold and directions on the tangent space T_pM .

Curve Representation of Tangent Vectors

Given a point $p \in M$ of a d -dimensional submanifold $M \subset \mathbb{R}^n$, we can represent a tangent vector $v \in T_p M$ as a curve $c : (-\varepsilon, \varepsilon) \rightarrow M$ with $c(0) = p$.

To see this, let us look at the manifold from the point of view of a coordinate mapping $x : U \rightarrow M$ with $0 \in U \subset \mathbb{R}^d$ and $x(0) = p$.

Since $v \in T_p M = \text{Im}(Dx(0))$, we know that there is an $h \in \mathbb{R}^d$ such that $Dx(0)[h] = v$.

Using

$$c : (-\varepsilon, \varepsilon) \rightarrow M$$

$$c(t) = x(t \cdot h),$$

we have

$$Dc(0) = Dx(0 \cdot h) \cdot h = v.$$

Alternative Definition of Tangent Vectors

Given a point $p \in M$ of a d -dimensional submanifold $M \subset \mathbb{R}^n$, we define

$$\mathcal{C}_p M := \{c: (-\varepsilon, \varepsilon) \rightarrow M \mid \exists \varepsilon > 0 : c \text{ is smooth and } c(0) = p\}.$$

The goal is to define $T_p M$ by defining an equivalence relation on $\mathcal{C}_p M$:

$$c_1 \sim c_2 \quad :\Leftrightarrow \quad Dc_1(0) = Dc_2(0),$$

It is easy to check that \sim satisfies *reflexivity*, *symmetry* and *transitivity*.

It turns out $T_p M = \mathcal{C}_p M / \sim$, which provides us with an alternative definition for the tangent space $T_p M$.

The advantage of this rather technical definition is that for any $v \in T_p M$ we can **choose a curve** $c \in v$ that passes through p and vice versa, *i.e.*, any curve c that passes through a point p defines a tangent vector $v := [c]$.

Differential as Push-Forward

Given two submanifolds M and N as well as a function $f: M \rightarrow N$. For $p \in M$ and $q = f(p) \in N$, the differential $Df(p)$ is the **push-forward**

$$\begin{aligned} Df(p): T_p M &\rightarrow T_q N \\ [c] &\mapsto [f \circ c] \end{aligned}$$

Assuming that $x_p: U_p \rightarrow M$ is a coordinate mapping for $p = x_p(0)$ and $x_q: U_q \rightarrow N$ is a coordinate mapping for $q = x_q(0)$, the push-forward definition becomes

$$Df(p)[v] = \left. \frac{\partial}{\partial t} (f \circ x_p)(t \cdot h) \right|_{t=0},$$

where $v = Dx_p(0)[h] = \left. \frac{\partial}{\partial t} x_p(t \cdot h) \right|_{t=0}$.

Push-Forward of Submanifolds

Given two submanifolds $M \subset \mathbb{R}^m$ and $N \subset \mathbb{R}^n$ as well as a function $f: \mathbb{R}^m \rightarrow \mathbb{R}^n$ with $f(M) = N$. For $p \in M$ and $q = f(p) \in N$, $Df(p)$ is

$$Df(p): T_p M \rightarrow T_q N \\ [c] \mapsto [f \circ c]$$

Assuming that $x_p: U_p \rightarrow M$ is a coordinate mapping for $p = x_p(0)$ and $x_q: U_q \rightarrow N$ is a coordinate mapping for $q = x_q(0)$, the push-forward definition becomes

$$Df(p)[v] = \left. \frac{\partial}{\partial t} (x_q \circ x_q^{-1} \circ f \circ x_p)(t \cdot h) \right|_{t=0},$$

where $v = Dx_p(0)[h]$.

It is easy to show that the push-forward is a linear mapping. Exercise.

Coordinate Interpretation

Given coordinate mappings $x_p: U_p \rightarrow M$ and $x_q: U_q \rightarrow N$ with $p = x_p(0)$ and $q = f(p) = x_q(0)$, the differential becomes

$$Df(p)[v] = \left. \frac{\partial}{\partial t} (x_q \circ x_q^{-1} \circ f \circ x_p)(t \cdot h) \right|_{t=0}.$$

If we were to apply the chain rule, we would obtain

$$Df(p)[v] = \mathbf{D}(x_q)(x^{-1}(q)) \cdot \mathbf{D}(x_q^{-1})(q) \cdot \mathbf{D}f(p) \cdot \mathbf{D}x_p(0) \cdot \mathbf{h}$$

- $Dx_p(0)h$ defines the tangent vector $v \in T_pM$.
- $Df(p)$ is the differential of f ignoring the submanifolds M and N .
- $Df(p)Dx_p(0)h$ is the differential of f only taking M into account.
- $D(x_q^{-1})(q)$ is the pseudo-inverse of $D(x_q)(x_q^{-1}(q))$.
- $D(x_q)(x^{-1}(q)) \cdot D(x_q^{-1})(q)$ projects a vector onto T_qN .

How to Compute the Differential

Given two submanifolds $M \subset \mathbb{R}^m$ and $N \subset \mathbb{R}^n$ as well as a function $f: \mathbb{R}^m \rightarrow \mathbb{R}^n$ with $f(M) = N$. For $p \in M$ and $q = f(p) \in N$, $Df(p)$ is

$$Df(p): T_p M \rightarrow T_q N \\ [c] \mapsto [f \circ c]$$

Assuming that $x_p: U_p \rightarrow M$ is a coordinate mapping for $p = x_p(0)$ and $x_q: U_q \rightarrow N$ is a coordinate mapping for $q = x_q(0)$, the push-forward definition becomes

$$Df(p)[v] = \pi_{T_q N}(D(f \circ x_p)(0)[h])$$

where $v = Dx_p(0)[h]$ and

$\pi_{T_q N}(v)$ is the orthogonal projection of $v \in \mathbb{R}^n$ to $T_q N$.

What is a Matrix? (Recap)

Linear mappings are commonly represented by matrices. We want to emphasize the difference between a matrix and a linear mapping.

Given an m -dimensional \mathbb{R} -vector space X , an n -dimensional \mathbb{R} -vector space Y and a linear mapping $L: X \rightarrow Y$, we can represent L by finite many scalars.

To this end, let $\mathcal{B}_X = \{x_1, \dots, x_m\}$ and $\mathcal{B}_Y = \{y_1, \dots, y_n\}$ bases of X and Y respectively. Then we know that for each $x_j \in \mathcal{B}_X$ we have

$$L(x_j) = \sum_{i=1}^n a_{ij} y_i.$$

for some $a_{ij} \in \mathbb{R}$.

We write this a_{ij} in a matrix A and call $A = \mathcal{M}_{\mathcal{B}_Y}^{\mathcal{B}_X}(L) \in \mathbb{R}^{n \times m}$ the representing matrix of L with respect to the basis \mathcal{B}_X and \mathcal{B}_Y .

Matrix of the Differential

Given two submanifolds M and N as well as a function $f: M \rightarrow N$. For $p \in M$ and $q = f(p) \in N$, the differential $Df(p): T_pM \rightarrow T_qN$ is a linear mapping, but in general we do not have a canonical matrix representation.

This means that any basis \mathcal{B}_p of T_pM and \mathcal{B}_q of T_qN would define a different matrix $\mathcal{M}_{\mathcal{B}_q}^{\mathcal{B}_p}(Df(p)) \in \mathbb{R}^{n \times m}$ with $n = \dim(N)$ and $m = \dim(M)$.

Since $T_pM = \text{Im}(Dx(0))$, $\mathcal{B}_p = \{Dx(0)[e_1], \dots, Dx(0)[e_m]\}$ would be a natural way of defining a basis of T_pM . Nonetheless, the resulting matrix would then depend on the coordinate mappings x_p and x_q that we choose for $p \in M$ and $q \in N$ respectively.

While there is no unique matrix that describes the differential, it is important to note that the image $\text{Im } Df(p)$ is independent of the chosen coordinate mappings. This was used for the definition of the tangent space.

Planar Curves and Normals

Given a 2D object O and its boundary, the 1D submanifold $M := \partial O$, we like to define the normal vector $n(p)$ for each point $p \in M$.

Given a coordinate mapping $x: U \rightarrow M$ with $x(0) = p$, we have $T_p M = \text{Im}(Dx(0))$ and a normal vector might be defined via

$$n(p) = \frac{1}{\|Dx(0)\|} \begin{pmatrix} +Dx^2(0) \\ -Dx^1(0) \end{pmatrix} \in \mathbb{S}^1$$

Since M is of codimension 1, $n(p)$ is up to the sign uniquely defined.

Thus, we have a smooth mapping

$$n: M \rightarrow \mathbb{S}^1$$

that defines a unique normal vector field of M . Why?

Differential of the Normal Mapping

Given a point $p \in M$ of $M \subset \mathbb{R}^2$, we have

$$Dn(p): T_p M \rightarrow T_{n(p)} \mathbb{S}^1.$$

Since we have

$$T_{n(p)} \mathbb{S}^1 = n(p)^\perp = T_p M,$$

we know that $Dn(p)$ is an **endomorphism**, i.e., a linear mapping that maps the vector space $T_p M$ onto itself.

Because $\dim T_p M = 1$, $Dn(p)$ maps a vector $v \in T_p M$ to $\kappa(p) \cdot v$.

This scalar value $\kappa(p) \in \mathbb{R}$ is called the **curvature** of M at the position p .

Derivative of the Normal Field

If we take the derivative of $N = (N^1, N^2): U \rightarrow \mathbb{R}^2$, $t \mapsto n \circ x(t)$, we obtain

$$\begin{aligned} \frac{\partial}{\partial t} N^1(x(t)) &= \frac{\partial}{\partial t} \frac{\dot{x}^2(t)}{\|\dot{x}(t)\|} = \frac{\ddot{x}^2(t) \|\dot{x}(t)\| - \dot{x}^2(t) \frac{\ddot{x}^1(t) + \ddot{x}^2(t)}{\|\dot{x}(t)\|}}{\|\dot{x}(t)\|^2} \\ &= \frac{\ddot{x}^2(t) \|\dot{x}(t)\|^2 - \dot{x}^2(t) \cdot (\ddot{x}^1(t) + \ddot{x}^2(t))}{\|\dot{x}(t)\|^3} \\ \frac{\partial}{\partial t} N^2(x(t)) &= \frac{-\ddot{x}^1(t) \|\dot{x}(t)\|^2 + \dot{x}^1(t) \cdot (\ddot{x}^1(t) + \ddot{x}^2(t))}{\|\dot{x}(t)\|^3} \end{aligned}$$

Note that $DN(p)$ is not necessarily in $T_p M$. Thus, we have to project it onto $T_p M$. To this end, let us choose $\{\dot{x}(t)\}$ as the base of $T_p M$.

Curvature

Overall, we have $Dn(p)[\dot{x}(t)] = \kappa(t) \cdot \dot{x}(t)$.

Therefore, we have

$$\begin{aligned}\kappa(t) &= \frac{\langle \dot{N}(t), \dot{x}(t) \rangle}{\|\dot{x}(t)\|^2} = \frac{\dot{x}^1(t)\ddot{x}^2(t)\|\dot{x}(t)\|^2 - \dot{x}^2(t)\ddot{x}^1(t)\|\dot{x}(t)\|^2}{\|\dot{x}(t)\|^5} \\ &= \frac{\det(\dot{x}(t), \ddot{x}(t))}{\|\dot{x}(t)\|^3}\end{aligned}$$

By construction, we know that curvature is invariant with respect to

- Translation. Why?
- Rotation. Why?
- Reparametrization. Why?

Curvature of Implicit Submanifolds

If $F: \mathbb{R}^2 \rightarrow \mathbb{R}$ has the regular value $c \in \mathbb{R}$, how can we use F in order to compute the curvature of M at $p \in M$?

The normal field n can be defined as $n(p) = \frac{\nabla F(p)}{\|\nabla F(p)\|}$.

Since n is also defined in a neighborhood of M , we can compute its derivative $Dn: M \rightarrow \mathbb{R}^{2 \times 2}$. If we write the linear mapping $Dn(p)$ with respect to the basis $\mathcal{B}_p = \{\nabla F(p), \nabla F(p)^\perp\}$, we obtain

$$\mathcal{M}_{\mathcal{B}_p}^{\mathcal{B}_p}(Dn(p)) = \begin{pmatrix} 0 & * \\ * & \kappa(p) \end{pmatrix}.$$

Therefore, we have

$$\kappa(p) = \operatorname{tr} Dn(p) = \operatorname{div} \left(\frac{\nabla F(p)}{\|\nabla F(p)\|} \right)$$

Literature

Differential

- Cauchy, *Cours d'Analyse de l'École Royale Polytechnique; I.re Partie. Analyse algébrique*, 1821.