

Chapter 1

Convex Analysis

Convex Optimization for Machine Learning & Computer Vision
SS 2018

Tao Wu
Emanuel Laude
Zhenzhang Ye

Computer Vision Group
Department of Informatics
TU Munich

Convex Analysis

Tao Wu
Emanuel Laude
Zhenzhang Ye



Convex Set

Convex Function

Existence of Minimizer



Convex Set

Notations

- \mathbb{E} is a Euclidean space (finite dimensional vector space), equipped with the inner product $\langle \cdot, \cdot \rangle$, e.g. $\langle u, v \rangle = u^\top v$.
- C is a closed, convex subset of \mathbb{E} .
- J is a convex objective function.

Convex optimization

$$\text{minimize } J(u) \quad \text{over } u \in C.$$

First questions:

- What is a convex set?
- What is a convex function?



Convex Set

Convex Function

Existence of Minimizer

Convex set


Definition

A set C is said to be **convex** if


$$\alpha u + (1 - \alpha)v \in C, \quad \forall u, v \in C, \quad \forall \alpha \in [0, 1].$$



convex



non-convex



Definition

- A set $C \subset \mathbb{E}$ is **open** if $\forall u \in C, \exists \epsilon > 0$ s.t. $B_\epsilon(u) \subset C$, where $B_\epsilon(u) := \{v \in \mathbb{E} : \|v - u\| < \epsilon\}$.
- A set $C \subset \mathbb{E}$ is **closed** if its complement $\mathbb{E} \setminus C$ is open.
- The **closure** of a set $C \subset \mathbb{E}$ is

$$\text{cl } C = \{u \in \mathbb{E} : \exists \{u^k\} \subset C \text{ s.t. } \lim_{k \rightarrow \infty} u^k = u\}.$$

- The **interior** of a set $C \subset \mathbb{E}$ is

$$\text{int } C = \{u \in C : \exists \epsilon > 0 \text{ s.t. } B_\epsilon(u) \subset C\}.$$





Definition

- A set $C \subset \mathbb{E}$ is **open** if $\forall u \in C, \exists \epsilon > 0$ s.t. $B_\epsilon(u) \subset C$, where $B_\epsilon(u) := \{v \in \mathbb{E} : \|v - u\| < \epsilon\}$.
- A set $C \subset \mathbb{E}$ is **closed** if its complement $\mathbb{E} \setminus C$ is open.
- The **closure** of a set $C \subset \mathbb{E}$ is

$$\text{cl } C = \{u \in \mathbb{E} : \exists \{u^k\} \subset C \text{ s.t. } \lim_{k \rightarrow \infty} u^k = u\}.$$

- The **interior** of a set $C \subset \mathbb{E}$ is

$$\text{int } C = \{u \in C : \exists \epsilon > 0 \text{ s.t. } B_\epsilon(u) \subset C\}.$$

- The **relative interior** of a set $C \subset \mathbb{E}$ is

$$\begin{aligned} \text{rint } C &:= \{u \in C : \exists \epsilon > 0 \text{ s.t. } B_\epsilon(u) \cap \text{aff } C \subset C\} \\ &= \{u \in C : \forall v \in C, \exists \alpha > 1 \text{ s.t. } v + \alpha(u - v) \in C\} \end{aligned}$$

if C is convex. Here $\text{aff } C$ stands for the **affine hull** of C .

The following operations preserve the convexity:

- Intersection: $C_1 \cap C_2$
- Summation: $C_1 + C_2 := \{u^1 + u^2 : u^1 \in C_1, u^2 \in C_2\}$
- Closure: $\text{cl } C$
- Interior: $\text{int } C$

– The union of convex sets is not convex in general.





The following operations preserve the convexity:

- Intersection: $C_1 \cap C_2$
- Summation: $C_1 + C_2 := \{u^1 + u^2 : u^1 \in C_1, u^2 \in C_2\}$
- Closure: $\text{cl } C$
- Interior: $\text{int } C$


– The union of convex sets is not convex in general.

– *Polyhedral sets* are always convex; *cones* are not necessarily convex.

Convex cone

C is a **cone** if $C = \alpha C$ for any $\alpha > 0$. C is a **convex cone** if C is a cone and is convex as well.

Separation of convex sets



Source: Wikipedia.

Theorem (separation of convex sets)

Let C_1, C_2 be nonempty convex subsets in \mathbb{E} s.t. $C_1 \cap C_2 = \emptyset$ and C_1 is open. Then there exists a hyperplane separating C_1 and C_2 , i.e. $\exists v \in \mathbb{E}, v \neq 0, \alpha \in \mathbb{R}$ s.t.

$$\langle v, u^1 \rangle \geq \alpha \geq \langle v, u^2 \rangle, \quad \forall u^1 \in C_1, u^2 \in C_2.$$

Proof: on board.



Separation of convex sets



Theorem (separation of convex sets)

Let C_1, C_2 be nonempty convex subsets in \mathbb{E} s.t. $C_1 \cap C_2 = \emptyset$ and C_1 is open. Then there exists a hyperplane separating C_1 and C_2 , i.e. $\exists v \in \mathbb{E}, v \neq 0, \alpha \in \mathbb{R}$ s.t.

$$\langle v, u^1 \rangle \geq \alpha \geq \langle v, u^2 \rangle, \quad \forall u^1 \in C_1, u^2 \in C_2.$$

Proof: on board.

Remarks

- 1 The proof works in any Hilbert space.
- 2 Corollary: In a Hilbert space, any (strongly) closed convex subset \overline{C} is weakly closed.
- 3 The above theorem generalizes to any topological vector space, known as the *Hahn-Banach theorem*.



Convex Function



- An **extended real-valued function** J maps from \mathbb{E} to $\bar{\mathbb{R}} := \mathbb{R} \cup \{\infty\}$.
- The **domain** of $J : \mathbb{E} \rightarrow \bar{\mathbb{R}}$ is

$$\text{dom } J = \{u \in \mathbb{E} : J(u) < \infty\}.$$

- The function $J : \mathbb{E} \rightarrow \bar{\mathbb{R}}$ is **proper** if $\text{dom } J \neq \emptyset$.

Definition

We say $J : \mathbb{E} \rightarrow \bar{\mathbb{R}}$ is a **convex function** if

- 1 $\text{dom } J$ is a convex set.
- 2 For all $u, v \in \text{dom } J$ and $\alpha \in [0, 1]$ it holds that

$$J(\alpha u + (1 - \alpha)v) \leq \alpha J(u) + (1 - \alpha)J(v).$$

We say J is **strictly convex** if the above inequality is strict for all $\alpha \in (0, 1)$ and $u \neq v$.

Examples

- $J_{data}(u) = \|u - f\|_q^q$ where $q \geq 1$ and $\|\cdot\|_q$ is the ℓ^q -norm.
- $J_{regu}(u) = \|Ku\|_p^p$ where K is linear transform and $p \geq 1$.
- $J(u) = J_{data}(u) + \alpha J_{regu}(u)$.



Examples

- $J_{data}(u) = \|u - f\|_q^q$ where $q \geq 1$ and $\|\cdot\|_q$ is the ℓ^q -norm.
- $J_{regu}(u) = \|Ku\|_p^p$ where K is linear transform and $p \geq 1$.
- $J(u) = J_{data}(u) + \alpha J_{regu}(u)$.
- (Binary) entropy: $J_\epsilon(u) = \epsilon(u \log(u) + (1 - u) \log(1 - u))$.
- Soft max: $J_\epsilon^*(v) = \epsilon \log(1 + \exp(v/\epsilon)) \approx \max(v, 0)$.





- $J_{data}(u) = \|u - f\|_q^q$ where $q \geq 1$ and $\|\cdot\|_q$ is the ℓ^q -norm.
- $J_{regu}(u) = \|Ku\|_p^p$ where K is linear transform and $p \geq 1$.
- $J(u) = J_{data}(u) + \alpha J_{regu}(u)$.
- (Binary) entropy: $J_\epsilon(u) = \epsilon(u \log(u) + (1 - u) \log(1 - u))$.
- Soft max: $J_\epsilon^*(v) = \epsilon \log(1 + \exp(v/\epsilon)) \approx \max(v, 0)$.
- **Indicator function** ($C \subset \mathbb{E}$ is closed and convex):

$$\delta_C(u) = \begin{cases} 0 & \text{if } u \in C, \\ \infty & \text{otherwise.} \end{cases}$$

- Formulate constrained optimization with indicator function:

$$\min J(u) \text{ over } u \in C. \Leftrightarrow \min J(u) + \delta_C(u) \text{ over } u \in \mathbb{E}.$$



(As exercises)

- Any norm (over a normed vector space) is a convex function.
- J is a convex function and K is a linear transform $\Rightarrow J(K\cdot)$ is convex function.
- (Jensen's inequality) $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ is convex iff

$$J\left(\sum_{i=1}^n \alpha_i u^i\right) \leq \sum_{i=1}^n \alpha_i J(u^i),$$


whenever $\{u^i\}_{i=1}^n \subset \mathbb{E}$, $\{\alpha_i\}_{i=1}^n \subset [0, 1]$, $\sum_{i=1}^n \alpha_i = 1$.

Epigraph

Definition

The **epigraph** of a proper function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ is

$$\text{epi } J = \{(u, \alpha) \in \mathbb{E} \times \mathbb{R} : J(u) \leq \alpha\}.$$



Theorem

A proper function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ is convex (resp. strictly convex) iff $\text{epi } J$ is a convex (resp. strictly convex) set.

Proof: as exercise.



Definition

Assume $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ with $\text{rint dom } J \neq \emptyset$. We say J is **locally Lipschitz** at $u \in \text{rint dom } J$ with modulus $L_u > 0$ if there exists $\epsilon > 0$ s.t.

$$|J(u^1) - J(u^2)| \leq L_u \|u^1 - u^2\| \quad \forall u^1, u^2 \in B_\epsilon(u) \cap \text{rint dom } J.$$



Convex Set

Convex Function

Existence of Minimizer



Definition

Assume $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ with $\text{rint dom } J \neq \emptyset$. We say J is **locally Lipschitz** at $u \in \text{rint dom } J$ with modulus $L_u > 0$ if there exists $\epsilon > 0$ s.t.

$$|J(u^1) - J(u^2)| \leq L_u \|u^1 - u^2\| \quad \forall u^1, u^2 \in B_\epsilon(u) \cap \text{rint dom } J.$$

Theorem

A proper convex function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ is locally Lipschitz at any $u \in \text{rint dom } J$.

Proof: on board.



Existence of Minimizer

Global vs. Local minimizer

Recall the optimization of $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$:

$$\text{minimize } J(u) \quad \text{over } u \in \mathbb{E}.$$



Definition

- 1 $u^* \in \mathbb{E}$ is a **global minimizer** if $J(u^*) \leq J(u)$ for all $u \in \mathbb{E}$.
- 2 u^* is a **local minimizer** if $\exists \epsilon > 0$ s.t. $J(u^*) \leq J(u)$ for all $u \in B_\epsilon(u^*)$.
- 3 In the above definitions, a global/local minimizer is **strict** if $J(u^*) \leq J(u)$ is replaced by $J(u^*) < J(u)$.

Global vs. Local minimizer

Recall the optimization of $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$:

$$\text{minimize } J(u) \quad \text{over } u \in \mathbb{E}.$$



Definition

- 1 $u^* \in \mathbb{E}$ is a **global minimizer** if $J(u^*) \leq J(u)$ for all $u \in \mathbb{E}$.
- 2 u^* is a **local minimizer** if $\exists \epsilon > 0$ s.t. $J(u^*) \leq J(u)$ for all $u \in B_\epsilon(u^*)$.
- 3 In the above definitions, a global/local minimizer is **strict** if $J(u^*) \leq J(u)$ is replaced by $J(u^*) < J(u)$.

Theorem

For any proper convex function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$, if $u^* \in \text{dom } J$ is a local minimizer of J , then it is also a global minimizer.

Proof: on board.

Does a minimizer always exist?




- Consider

$$\text{minimize } J(u) \quad \text{over } u \in \mathbb{E},$$

where $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ is a proper, convex function.

- Some counterexamples for $J : \mathbb{R} \rightarrow \overline{\mathbb{R}}$:




- We shall formalize our observations and derive sufficient conditions for existence.

Sufficient conditions for existence

Definition

- 1 J is **bounded from below** if $J(\cdot) \geq C$ for some $C \in \mathbb{R}$.
- 2 J is **coercive** if $J(u) \rightarrow \infty$ whenever $\|u\| \rightarrow \infty$.
 - Proposition: J is coercive if $\text{dom } J$ is bounded.
- 3 J is **lower semi-continuous** (lsc) at u^* if

$$J(u^*) \leq \liminf_{u \rightarrow u^*} J(u).$$



- Proposition: J is lsc iff $\text{epi } J$ is closed.



Sufficient conditions for existence

Definition

- 1 J is **bounded from below** if $J(\cdot) \geq C$ for some $C \in \mathbb{R}$.
- 2 J is **coercive** if $J(u) \rightarrow \infty$ whenever $\|u\| \rightarrow \infty$.
- 3 J is **lower semi-continuous** (lsc) at u^* if

$$J(u^*) \leq \liminf_{u \rightarrow u^*} J(u).$$



Theorem

Any proper function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$, which is bounded from below, coercive, and lsc (everywhere), has a (global) minimizer.

Proof: on board.

Sufficient conditions for existence

Definition

- 1 J is **bounded from below** if $J(\cdot) \geq C$ for some $C \in \mathbb{R}$.
- 2 J is **coercive** if $J(u) \rightarrow \infty$ whenever $\|u\| \rightarrow \infty$.
- 3 J is **lower semi-continuous** (lsc) at u^* if

$$J(u^*) \leq \liminf_{u \rightarrow u^*} J(u).$$



Theorem

Any proper function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$, which is bounded from below, coercive, and lsc (everywhere), has a (global) minimizer.

Proof: on board.

Remarks for infinite dimensions

- 1 Weak compactness in reflexive Banach (e.g. Hilbert) sp.
- 2 J is convex and strongly lsc $\Rightarrow J$ is weakly lsc.



- Recall that a function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ is strictly convex if

$$J(\alpha u + (1 - \alpha)v) < \alpha J(u) + (1 - \alpha)J(v),$$

for all $u, v \in \text{dom } J$, $u \neq v$, $\alpha \in (0, 1)$.

Theorem

The minimizer of a strictly convex function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ is unique.

Proof: on board.