

# Chapter 1

## Convex Analysis

*Convex Optimization for Machine Learning & Computer Vision*  
SS 2018

Tao Wu  
Emanuel Laude  
Zhenzhang Ye

Computer Vision Group  
Department of Informatics  
TU Munich

Convex Analysis

Tao Wu  
Emanuel Laude  
Zhenzhang Ye



Convex Set

Convex Function

Existence of Minimizer

Subdifferential



# Convex Set



## Notations

- $\mathbb{E}$  is a Euclidean space (finite dimensional vector space), equipped with the inner product  $\langle \cdot, \cdot \rangle$ , e.g.  $\langle u, v \rangle = u^\top v$ .
- $C$  is a closed, convex subset of  $\mathbb{E}$ .
- $J$  is a convex objective function.

## Convex Set

## Convex Function

## Existence of Minimizer

## Subdifferential

## Convex optimization

$$\text{minimize } J(u) \quad \text{over } u \in C.$$

First questions:


- What is a convex set?
- What is a convex function?

## Definition


A set  $C$  is said to be **convex** if

$$\alpha u + (1 - \alpha)v \in C, \quad \forall u, v \in C, \quad \forall \alpha \in [0, 1].$$

convex



non-convex



### Definition

- A set  $C \subset \mathbb{E}$  is **open** if  $\forall u \in C, \exists \epsilon > 0$  s.t.  $B_\epsilon(u) \subset C$ , where  $B_\epsilon(u) := \{v \in \mathbb{E} : \|v - u\| < \epsilon\}$ .
- A set  $C \subset \mathbb{E}$  is **closed** if its complement  $\mathbb{E} \setminus C$  is open.
- The **closure** of a set  $C \subset \mathbb{E}$  is

$$\text{cl } C = \{u \in \mathbb{E} : \exists \{u^k\} \subset C \text{ s.t. } \lim_{k \rightarrow \infty} u^k = u\}.$$

- The **interior** of a set  $C \subset \mathbb{E}$  is

$$\text{int } C = \{u \in C : \exists \epsilon > 0 \text{ s.t. } B_\epsilon(u) \subset C\}.$$





### Definition

- A set  $C \subset \mathbb{E}$  is **open** if  $\forall u \in C, \exists \epsilon > 0$  s.t.  $B_\epsilon(u) \subset C$ , where  $B_\epsilon(u) := \{v \in \mathbb{E} : \|v - u\| < \epsilon\}$ .
- A set  $C \subset \mathbb{E}$  is **closed** if its complement  $\mathbb{E} \setminus C$  is open.
- The **closure** of a set  $C \subset \mathbb{E}$  is

$$\text{cl } C = \{u \in \mathbb{E} : \exists \{u^k\} \subset C \text{ s.t. } \lim_{k \rightarrow \infty} u^k = u\}.$$

- The **interior** of a set  $C \subset \mathbb{E}$  is

$$\text{int } C = \{u \in C : \exists \epsilon > 0 \text{ s.t. } B_\epsilon(u) \subset C\}.$$

- The **relative interior** of a set  $C \subset \mathbb{E}$  is

$$\begin{aligned} \text{rint } C &:= \{u \in C : \exists \epsilon > 0 \text{ s.t. } B_\epsilon(u) \cap \text{aff } C \subset C\} \\ &= \{u \in C : \forall v \in C, \exists \alpha > 1 \text{ s.t. } v + \alpha(u - v) \in C\} \end{aligned}$$

if  $C$  is convex. Here  $\text{aff } C$  stands for the **affine hull** of  $C$ .



The following operations preserve the convexity:

- Intersection:  $C_1 \cap C_2$
- Summation:  $C_1 + C_2 := \{u^1 + u^2 : u^1 \in C_1, u^2 \in C_2\}$
- Closure:  $\text{cl } C$
- Interior:  $\text{int } C$

– The union of convex sets is not convex in general.



The following operations preserve the convexity:

- Intersection:  $C_1 \cap C_2$
- Summation:  $C_1 + C_2 := \{u^1 + u^2 : u^1 \in C_1, u^2 \in C_2\}$
- Closure:  $\text{cl } C$
- Interior:  $\text{int } C$

– The union of convex sets is not convex in general.


– *Polyhedral sets* are always convex; *cones* are not necessarily convex.

### Convex cone

$C$  is a **cone** if  $C = \alpha C$  for any  $\alpha > 0$ .  $C$  is a **convex cone** if  $C$  is a cone and is convex as well.



## Separation of convex sets



Source: Wikipedia.

### Theorem (separation of convex sets)

Let  $C_1, C_2$  be nonempty convex subsets in  $\mathbb{E}$  s.t.  $C_1 \cap C_2 = \emptyset$  and  $C_1$  is open. Then there exists a hyperplane separating  $C_1$  and  $C_2$ , i.e.  $\exists v \in \mathbb{E}, v \neq 0, \alpha \in \mathbb{R}$  s.t.

$$\langle v, u^1 \rangle \geq \alpha \geq \langle v, u^2 \rangle, \quad \forall u^1 \in C_1, u^2 \in C_2.$$

Proof: on board.



# Separation of convex sets



## Theorem (separation of convex sets)

Let  $C_1, C_2$  be nonempty convex subsets in  $\mathbb{E}$  s.t.  $C_1 \cap C_2 = \emptyset$  and  $C_1$  is open. Then there exists a hyperplane separating  $C_1$  and  $C_2$ , i.e.  $\exists v \in \mathbb{E}, v \neq 0, \alpha \in \mathbb{R}$  s.t.

$$\langle v, u^1 \rangle \geq \alpha \geq \langle v, u^2 \rangle, \quad \forall u^1 \in C_1, u^2 \in C_2.$$

Proof: on board.

## Remarks

- 1 The proof works in any Hilbert space.
- 2 Corollary: In a Hilbert space, any (strongly) closed convex subset  $\overline{C}$  is weakly closed.
- 3 The above theorem generalizes to any topological vector space, known as the *Hahn-Banach theorem*.



# Convex Function



- An **extended real-valued function**  $J$  maps from  $\mathbb{E}$  to  $\bar{\mathbb{R}} := \mathbb{R} \cup \{\infty\}$ .
- The **domain** of  $J : \mathbb{E} \rightarrow \bar{\mathbb{R}}$  is

$$\text{dom } J = \{u \in \mathbb{E} : J(u) < \infty\}.$$

- The function  $J : \mathbb{E} \rightarrow \bar{\mathbb{R}}$  is **proper** if  $\text{dom } J \neq \emptyset$ .

## Definition

We say  $J : \mathbb{E} \rightarrow \bar{\mathbb{R}}$  is a **convex function** if

- 1  $\text{dom } J$  is a convex set.
- 2 For all  $u, v \in \text{dom } J$  and  $\alpha \in [0, 1]$  it holds that

$$J(\alpha u + (1 - \alpha)v) \leq \alpha J(u) + (1 - \alpha)J(v).$$

We say  $J$  is **strictly convex** if the above inequality is strict for all  $\alpha \in (0, 1)$  and  $u \neq v$ .

## Examples

- $J_{data}(u) = \|u - f\|_q^q$  where  $q \geq 1$  and  $\|\cdot\|_q$  is the  $\ell^q$ -norm.
- $J_{regu}(u) = \|Ku\|_p^p$  where  $K$  is linear transform and  $p \geq 1$ .
- $J(u) = J_{data}(u) + \alpha J_{regu}(u)$ .



## Examples

- $J_{data}(u) = \|u - f\|_q^q$  where  $q \geq 1$  and  $\|\cdot\|_q$  is the  $\ell^q$ -norm.
- $J_{regu}(u) = \|Ku\|_p^p$  where  $K$  is linear transform and  $p \geq 1$ .
- $J(u) = J_{data}(u) + \alpha J_{regu}(u)$ .
- (Binary) entropy:  $J_\epsilon(u) = \epsilon(u \log(u) + (1 - u) \log(1 - u))$ .
- Soft max:  $J_\epsilon^*(v) = \epsilon \log(1 + \exp(v/\epsilon)) \approx \max(v, 0)$ .





- $J_{data}(u) = \|u - f\|_q^q$  where  $q \geq 1$  and  $\|\cdot\|_q$  is the  $\ell^q$ -norm.
- $J_{regu}(u) = \|Ku\|_p^p$  where  $K$  is linear transform and  $p \geq 1$ .
- $J(u) = J_{data}(u) + \alpha J_{regu}(u)$ .
- (Binary) entropy:  $J_\epsilon(u) = \epsilon(u \log(u) + (1 - u) \log(1 - u))$ .
- Soft max:  $J_\epsilon^*(v) = \epsilon \log(1 + \exp(v/\epsilon)) \approx \max(v, 0)$ .
- **Indicator function** ( $C \subset \mathbb{E}$  is closed and convex):

$$\delta_C(u) = \begin{cases} 0 & \text{if } u \in C, \\ \infty & \text{otherwise.} \end{cases}$$

- Formulate constrained optimization with indicator function:

$$\min J(u) \text{ over } u \in C. \Leftrightarrow \min J(u) + \delta_C(u) \text{ over } u \in \mathbb{E}.$$



(As exercises)

- Any norm (over a normed vector space) is a convex function.
- $J$  is a convex function and  $K$  is a linear transform  $\Rightarrow J(K\cdot)$  is convex function.
- (Jensen's inequality)  $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$  is convex iff

$$J\left(\sum_{i=1}^n \alpha_i u^i\right) \leq \sum_{i=1}^n \alpha_i J(u^i),$$

whenever  $\{u^i\}_{i=1}^n \subset \mathbb{E}$ ,  $\{\alpha_i\}_{i=1}^n \subset [0, 1]$ ,  $\sum_{i=1}^n \alpha_i = 1$ .




# Epigraph

## Definition

The **epigraph** of a proper function  $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$  is

$$\text{epi } J = \{(u, \alpha) \in \mathbb{E} \times \mathbb{R} : J(u) \leq \alpha\}.$$



## Theorem

A proper function  $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$  is convex (resp. strictly convex) iff  $\text{epi } J$  is a convex (resp. strictly convex) set.

Proof: as exercise.



## Definition

Assume  $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$  with  $\text{rint dom } J \neq \emptyset$ . We say  $J$  is **locally Lipschitz** at  $u \in \text{rint dom } J$  with modulus  $L_u > 0$  if there exists  $\epsilon > 0$  s.t.

$$|J(u^1) - J(u^2)| \leq L_u \|u^1 - u^2\| \quad \forall u^1, u^2 \in B_\epsilon(u) \cap \text{rint dom } J.$$



Convex Set

Convex Function

Existence of Minimizer

Subdifferential



## Definition

Assume  $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$  with  $\text{rint dom } J \neq \emptyset$ . We say  $J$  is **locally Lipschitz** at  $u \in \text{rint dom } J$  with modulus  $L_u > 0$  if there exists  $\epsilon > 0$  s.t.

$$|J(u^1) - J(u^2)| \leq L_u \|u^1 - u^2\| \quad \forall u^1, u^2 \in B_\epsilon(u) \cap \text{rint dom } J.$$

## Theorem

A proper convex function  $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$  is locally Lipschitz at any  $u \in \text{rint dom } J$ .

Proof: on board.

Convex Set

Convex Function

Existence of Minimizer

Subdifferential



# Existence of Minimizer

Convex Set

Convex Function

Existence of Minimizer

Subdifferential

## Global vs. Local minimizer

Recall the optimization of  $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ :

$$\text{minimize } J(u) \quad \text{over } u \in \mathbb{E}.$$

### Definition

- 1  $u^* \in \mathbb{E}$  is a **global minimizer** if  $J(u^*) \leq J(u)$  for all  $u \in \mathbb{E}$ .
- 2  $u^*$  is a **local minimizer** if  $\exists \epsilon > 0$  s.t.  $J(u^*) \leq J(u)$  for all  $u \in B_\epsilon(u^*)$ .
- 3 In the above definitions, a global/local minimizer is **strict** if  $J(u^*) \leq J(u)$  is replaced by  $J(u^*) < J(u)$ .



## Global vs. Local minimizer

Recall the optimization of  $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ :

$$\text{minimize } J(u) \quad \text{over } u \in \mathbb{E}.$$

### Definition

- 1  $u^* \in \mathbb{E}$  is a **global minimizer** if  $J(u^*) \leq J(u)$  for all  $u \in \mathbb{E}$ .
- 2  $u^*$  is a **local minimizer** if  $\exists \epsilon > 0$  s.t.  $J(u^*) \leq J(u)$  for all  $u \in B_\epsilon(u^*)$ .
- 3 In the above definitions, a global/local minimizer is **strict** if  $J(u^*) \leq J(u)$  is replaced by  $J(u^*) < J(u)$ .

### Theorem

For any proper convex function  $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ , if  $u^* \in \text{dom } J$  is a local minimizer of  $J$ , then it is also a global minimizer.

Proof: on board.



# Does a minimizer always exist?




- Consider

$$\text{minimize } J(u) \quad \text{over } u \in \mathbb{E},$$

where  $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$  is a proper, convex function.

- Some counterexamples for  $J : \mathbb{R} \rightarrow \overline{\mathbb{R}}$ :





- We shall formalize our observations and derive sufficient conditions for existence.

# Sufficient conditions for existence

## Definition

- 1  $J$  is **bounded from below** if  $J(\cdot) \geq C$  for some  $C \in \mathbb{R}$ .
- 2  $J$  is **coercive** if  $J(u) \rightarrow \infty$  whenever  $\|u\| \rightarrow \infty$ .
  - Proposition:  $J$  is coercive if  $\text{dom } J$  is bounded.
- 3  $J$  is **lower semi-continuous** (lsc) at  $u^*$  if

$$J(u^*) \leq \liminf_{u \rightarrow u^*} J(u).$$



- Proposition:  $J$  is lsc iff  $\text{epi } J$  is closed.





# Sufficient conditions for existence

## Definition

- 1  $J$  is **bounded from below** if  $J(\cdot) \geq C$  for some  $C \in \mathbb{R}$ .
- 2  $J$  is **coercive** if  $J(u) \rightarrow \infty$  whenever  $\|u\| \rightarrow \infty$ .
- 3  $J$  is **lower semi-continuous** (lsc) at  $u^*$  if

$$J(u^*) \leq \liminf_{u \rightarrow u^*} J(u).$$

## Theorem

Any proper function  $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ , which is bounded from below, coercive, and lsc (everywhere), has a (global) minimizer.

Proof: on board.



## Sufficient conditions for existence

### Definition

- 1  $J$  is **bounded from below** if  $J(\cdot) \geq C$  for some  $C \in \mathbb{R}$ .
- 2  $J$  is **coercive** if  $J(u) \rightarrow \infty$  whenever  $\|u\| \rightarrow \infty$ .
- 3  $J$  is **lower semi-continuous** (lsc) at  $u^*$  if

$$J(u^*) \leq \liminf_{u \rightarrow u^*} J(u).$$



### Theorem

Any proper function  $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ , which is bounded from below, coercive, and lsc (everywhere), has a (global) minimizer.

Proof: on board.

### Remarks for infinite dimensions

- 1 Weak compactness in reflexive Banach (e.g. Hilbert) sp.
- 2  $J$  is convex and strongly lsc  $\Rightarrow J$  is weakly lsc.



- Recall that a function  $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$  is strictly convex if

$$J(\alpha u + (1 - \alpha)v) < \alpha J(u) + (1 - \alpha)J(v),$$

for all  $u, v \in \text{dom } J$ ,  $u \neq v$ ,  $\alpha \in (0, 1)$ .


## Theorem

The minimizer of a strictly convex function  $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$  is unique.

Proof: on board.



# Subdifferential




## Definition

$J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$  is called (Fréchet) **differentiable** at  $u \in \text{int dom } J$  and  $\nabla J(u) \in \mathbb{E}$  is the (Fréchet) **differential** of  $J$  at  $u$  if

$$\lim_{h \rightarrow 0} \frac{|J(u+h) - J(u) - \langle \nabla J(u), h \rangle|}{\|h\|} = 0.$$

$J$  is **continuously differentiable** at  $u \in \text{int dom } J$  if  $\nabla J(\cdot)$  is continuous on  $(\text{dom } J) \cap B_\epsilon(u)$  for some  $\epsilon > 0$ .





## Definition

$J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$  is called (Fréchet) **differentiable** at  $u \in \text{int dom } J$  and  $\nabla J(u) \in \mathbb{E}$  is the (Fréchet) **differential** of  $J$  at  $u$  if

$$\lim_{h \rightarrow 0} \frac{|J(u+h) - J(u) - \langle \nabla J(u), h \rangle|}{\|h\|} = 0.$$

$J$  is **continuously differentiable** at  $u \in \text{int dom } J$  if  $\nabla J(\cdot)$  is continuous on  $(\text{dom } J) \cap B_\epsilon(u)$  for some  $\epsilon > 0$ .


## Remark

If  $\mathbb{E}$  is a topological vector space,  $\nabla J(u)$  is treated as a *dual* object in  $\mathbb{E}^*$ , and  $\langle \nabla J(u), h \rangle_{\mathbb{E}^*, \mathbb{E}}$  as *duality pairing*.



## Subdifferential

Now we generalize differentiability from differentiable functions to non-differentiable convex functions.



### Definition


The **subdifferential** of a convex function  $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$  at  $u \in \text{dom } J$  is defined by

$$\partial J(u) = \{p \in \mathbb{E} : J(v) \geq J(u) + \langle p, v - u \rangle \quad \forall v \in \mathbb{E}\}.$$



## Subdifferential

Now we generalize differentiability from differentiable functions to non-differentiable convex functions.



### Definition

The **subdifferential** of a convex function  $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$  at  $u \in \text{dom } J$  is defined by

$$\partial J(u) = \{p \in \mathbb{E} : J(v) \geq J(u) + \langle p, v - u \rangle \quad \forall v \in \mathbb{E}\}.$$

### Geometric interpretation

$p \in \partial J(u)$  iff  $(p, -1)$  is a normal vector for the supporting hyperplane of  $\text{epi } J$  at  $(u, J(u))$ .





# Examples and basic facts

## Basic facts

- 1  $\partial J(\cdot)$  is a *set-valued* map.
- 2 If  $J$  is cont. differentiable at  $u$ , then  $\partial J(u) = \{\nabla J(u)\}$ .



# Examples and basic facts

## Basic facts

- 1  $\partial J(\cdot)$  is a *set-valued* map.
- 2 If  $J$  is cont. differentiable at  $u$ , then  $\partial J(u) = \{\nabla J(u)\}$ .

## Examples

- 1  $J(u) = \|u\| \Rightarrow \partial J(0) = \{p : \|u\|_* \leq 1\}$ , where  $\|\cdot\|_*$  is the **dual norm** of  $\|\cdot\|$ , i.e.,  $\|p\|_* = \sup\{\langle p, u \rangle : \|u\| \leq 1\}$ .
- 2 Given any closed, convex subset  $C \subset \mathbb{E}$  and  $u \in C$ ,

$$\partial \delta_C(u) = \{p \in \mathbb{E} : \langle p, v - u \rangle \leq 0 \forall v \in C\} =: N_C(u),$$

known as the **normal cone** of  $C$  at  $u$ .



### Basic facts

- 1  $\partial J(\cdot)$  is a *set-valued* map.
- 2 If  $J$  is cont. differentiable at  $u$ , then  $\partial J(u) = \{\nabla J(u)\}$ .



Convex Set

Convex Function

Existence of Minimizer

Subdifferential

### Examples

- 1  $J(u) = \|u\| \Rightarrow \partial J(0) = \{p : \|u\|_* \leq 1\}$ , where  $\|\cdot\|_*$  is the **dual norm** of  $\|\cdot\|$ , i.e.,  $\|p\|_* = \sup\{\langle p, u \rangle : \|u\| \leq 1\}$ .

- 2 Given any closed, convex subset  $C \subset \mathbb{E}$  and  $u \in C$ ,

$$\partial \delta_C(u) = \{p \in \mathbb{E} : \langle p, v - u \rangle \leq 0 \forall v \in C\} =: N_C(u),$$

known as the **normal cone** of  $C$  at  $u$ .

- 3 (Exercise)  $X \in \mathbb{R}^{m \times n} \mapsto \|X\|_{1,2} = \sum_{i=1}^m \left( \sum_{j=1}^n |X_{i,j}|^2 \right)^{1/2}$ .

- 4 (Exercise)  $X \in \mathbb{R}^{m \times n} \mapsto \|X\|_{nuclear} = \sum_i \sigma_i(X)$ , i.e., sum of singular values.

## Theorem (chain rule under linear transform)

Let  $\tilde{J}(\cdot) = J(K\cdot)$  with some convex function  $J$  and linear transform  $K$ . Then

$$\partial\tilde{J}(u) = K^\top \partial J(Ku)$$

whenever  $Ku \in \text{dom } J$ .

Example:  $J(u) = \|Ku\| \Rightarrow \partial J(u) = K^\top \partial \|\cdot\| \cdot \|(Ku)$ .



Convex Set

Convex Function

Existence of Minimizer

Subdifferential



## Theorem (chain rule under linear transform)

Let  $\tilde{J}(\cdot) = J(K\cdot)$  with some convex function  $J$  and linear transform  $K$ . Then

$$\partial\tilde{J}(u) = K^\top \partial J(Ku)$$

whenever  $Ku \in \text{dom } J$ .

Example:  $J(u) = \|Ku\| \Rightarrow \partial J(u) = K^\top \partial \|\cdot\|(Ku)$ .

## Theorem (summation rule)

If  $\tilde{J}(\cdot) = J_1(\cdot) + J_2(\cdot)$  for some convex functions  $J_1$  and  $J_2$ , then

$$\partial\tilde{J}(u) = \partial J_1(u) + \partial J_2(u)$$

for any  $u \in \text{dom } J_1 \cap \text{dom } J_2$ .

Warning: not true if  $J_1$  or  $J_2$  is non-convex, e.g.  $0 = |\cdot| + (-|\cdot|)$ .

## Properties of subdifferential map

### Theorem

Let  $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$  be a convex function. Then  $\partial J$  is a **monotone operator**, i.e.  $\forall u^1, u^2 \in \text{dom } J, p^1 \in \partial J(u^1), p^2 \in \partial J(u^2)$  :

$$\langle p^1 - p^2, u^1 - u^2 \rangle \geq 0.$$

Proof: on board.



## Properties of subdifferential map

### Theorem

Let  $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$  be a convex function. Then  $\partial J$  is a **monotone operator**, i.e.  $\forall u^1, u^2 \in \text{dom } J, p^1 \in \partial J(u^1), p^2 \in \partial J(u^2)$  :

$$\langle p^1 - p^2, u^1 - u^2 \rangle \geq 0.$$

Proof: on board.

### Theorem

Let  $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$  be a convex function. Then for any  $u \in \text{int dom } J$ ,  $\partial J(u)$  is a nonempty, compact, and convex subset.

Proof: on board.



## Properties of subdifferential map

### Theorem

Let  $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$  be a convex function. Then  $\partial J$  is a **monotone operator**, i.e.  $\forall u^1, u^2 \in \text{dom } J, p^1 \in \partial J(u^1), p^2 \in \partial J(u^2)$  :

$$\langle p^1 - p^2, u^1 - u^2 \rangle \geq 0.$$

Proof: on board.

### Theorem

Let  $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$  be a convex function. Then for any  $u \in \text{int dom } J$ ,  $\partial J(u)$  is a nonempty, compact, and convex subset.

Proof: on board.

### Theorem

Let  $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$  be a proper, convex, lsc function. Then the set-valued map  $\partial J(\cdot)$  is **closed**, i.e.  $p^* \in \partial J(u^*)$  whenever

$$\exists (u^k, p^k) \rightarrow (u^*, p^*) \in (\text{dom } J) \times \mathbb{E} \text{ s.t. } p^k \in \partial J(u^k) \quad \forall k.$$

Proof: on board.





# Optimality condition

## Theorem

Given any proper convex function  $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ , the sufficient and necessary condition for  $u^*$  being a (global) minimizer for  $J$  is

$$0 \in \partial J(u^*).$$

Proof: on board.



## Optimality condition

### Theorem

Given any proper convex function  $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ , the sufficient and necessary condition for  $u^*$  being a (global) minimizer for  $J$  is


$$0 \in \partial J(u^*).$$

Proof: on board.

### Constrained optimization as special case

If  $u^*$  minimizes  $\tilde{J} = J + \delta_C$  with convex function  $J : \mathbb{E} \rightarrow \mathbb{R}$  and closed convex subset  $C \in \mathbb{E}$ , then  $0 \in \partial \tilde{J}(u^*) \Leftrightarrow$

$$0 \in \partial J(u^*) + N_C(u^*).$$




# Optimality condition

## Constrained optimization as special case

If  $u^*$  minimizes  $\tilde{J} = J + \delta_C$  with convex function  $J : \mathbb{E} \rightarrow \mathbb{R}$  and closed convex subset  $C \in \mathbb{E}$ , then  $0 \in \partial \tilde{J}(u^*) \Leftrightarrow$

$$0 \in \partial J(u^*) + N_C(u^*).$$



## Remark

The optimality condition  $0 \in \partial J(u^*) + N_C(u^*)$  is *geometric*. More explicit characterization relies on the *algebraic* representation of  $N_C(u^*)$ , e.g., the **Karush-Kuhn-Tucker (KKT) conditions**, typically under certain *constraint qualifications*.



# Optimality condition

## Constrained optimization as special case

If  $u^*$  minimizes  $\tilde{J} = J + \delta_C$  with convex function  $J : \mathbb{E} \rightarrow \mathbb{R}$  and closed convex subset  $C \in \mathbb{E}$ , then  $0 \in \partial \tilde{J}(u^*) \Leftrightarrow$

$$0 \in \partial J(u^*) + N_C(u^*).$$

## Remark

The optimality condition  $0 \in \partial J(u^*) + N_C(u^*)$  is *geometric*. More explicit characterization relies on the *algebraic* representation of  $N_C(u^*)$ , e.g., the **Karush-Kuhn-Tucker (KKT) conditions**, typically under certain *constraint qualifications*.

## Example: Linear-inequality constraints

Let  $C = \{u \in \mathbb{R}^n : Au \leq b\}$  where  $b \in \mathbb{R}^m$ ,  $A \in \mathbb{R}^{m \times n}$  has linearly independent rows. Then

$$N_C(u) = \{A^\top \lambda : \lambda \geq 0, \lambda_i = 0 \text{ if } (Au - b)_i < 0\}.$$

