



Chapter 2

Optimization Algorithms

Convex Optimization for Machine Learning & Computer Vision
SS 2018

Tao Wu
Emanuel Laude
Zhenzhang Ye

Computer Vision Group
Department of Informatics
TU Munich



Gradient-based Methods

Unconstrained, differentiable, possibly nonconvex optimization

Problem setting:

$$\text{minimize } J(u) \quad \text{over } u \in \mathbb{E}.$$

Assume:

- 1 $J : \mathbb{E} \rightarrow \mathbb{R}$ is continuously differentiable.
- 2 There exists a global minimizer u^* . (Typically, an optimization algorithm seeks for a local minimizer s.t. $\nabla J(u^*) = 0$.)



Unconstrained, differentiable, possibly nonconvex optimization

Problem setting:

$$\text{minimize } J(u) \quad \text{over } u \in \mathbb{E}.$$

Assume:

- 1 $J : \mathbb{E} \rightarrow \mathbb{R}$ is continuously differentiable.
- 2 There exists a global minimizer u^* . (Typically, an optimization algorithm seeks for a local minimizer s.t. $\nabla J(u^*) = 0$.)

Methods under consideration:

- 1 (Scaled) gradient descent.
- 2 Line search method.
- 3 Majorize-minimize method.



Unconstrained, differentiable, possibly nonconvex optimization

Problem setting:

$$\text{minimize } J(u) \quad \text{over } u \in \mathbb{E}.$$

Assume:

- 1 $J : \mathbb{E} \rightarrow \mathbb{R}$ is continuously differentiable.
- 2 There exists a global minimizer u^* . (Typically, an optimization algorithm seeks for a local minimizer s.t. $\nabla J(u^*) = 0$.)


Methods under consideration:

- 1 (Scaled) gradient descent.
- 2 Line search method.
- 3 Majorize-minimize method.

Analytical questions:

- 1 Convergence (or not); global vs. local convergence.
- 2 Convergence rate (in special cases).





Descent method

Initialize $u^0 \in \mathbb{E}$. Iterate with $k = 0, 1, 2, \dots$

- 1 If the stopping criteria $\|\nabla J(u^k)\| \leq \epsilon$ is *not* satisfied, then continue; otherwise return u^k and stop.
- 2 Choose a **descent direction** $d^k \in \mathbb{E}$ s.t.

$$\langle \nabla J(u^k), d^k \rangle < 0.$$

- 3 Choose an “appropriate” step size $\tau^k > 0$, and update

$$u^{k+1} = u^k + \tau^k d^k.$$

Theorem

If $\langle \nabla J(u^k), d^k \rangle < 0$, then $J(u^k + \tau d^k) < J(u^k)$ for all sufficiently small $\tau > 0$.



Theorem

If $\langle \nabla J(u^k), d^k \rangle < 0$, then $J(u^k + \tau d^k) < J(u^k)$ for all sufficiently small $\tau > 0$.

Proof: Use the Taylor expansion:

$$\begin{aligned} J(u^k + \tau d^k) &= J(u^k) + \tau \langle \nabla J(u^k), d^k \rangle + o(\tau) \\ &= J(u^k) + \tau \left(\langle \nabla J(u^k), d^k \rangle + o(1) \right) < J(u^k) \quad \text{as } \tau \rightarrow 0^+. \end{aligned}$$



Theorem

If $\langle \nabla J(u^k), d^k \rangle < 0$, then $J(u^k + \tau d^k) < J(u^k)$ for all sufficiently small $\tau > 0$.

Proof: Use the Taylor expansion:


$$\begin{aligned} J(u^k + \tau d^k) &= J(u^k) + \tau \langle \nabla J(u^k), d^k \rangle + o(\tau) \\ &= J(u^k) + \tau \left(\langle \nabla J(u^k), d^k \rangle + o(1) \right) < J(u^k) \quad \text{as } \tau \rightarrow 0^+. \end{aligned}$$

Choices of descent direction

- 1 Scaled gradient: $d^k = -(H^k)^{-1} \nabla J(u^k)$.
- 2 Gradient/Steepest descent: $H^k = I$.
- 3 Newton: $H^k = \nabla^2 J(u^k)$, assuming J is twice continuously differentiable and $\nabla^2 J(u^k) \succ 0$.
- 4 Quasi-Newton: $H^k \approx \nabla^2 J(u^k)$, H^k is spd.



Gradient descent with exact line search




- Gradient descent with *exact* line search:

$$u^{k+1} = u^k - \tau^k \nabla J(u^k),$$

$$\tau^k = \arg \min_{\tau} J(u^k - \tau \nabla J(u^k)).$$



Gradient descent with exact line search




- Gradient descent with *exact* line search:

$$u^{k+1} = u^k - \tau^k \nabla J(u^k),$$
$$\tau^k = \arg \min_{\tau} J(u^k - \tau \nabla J(u^k)).$$

- Special case: $J(u) = \frac{1}{2} \langle u, Qu \rangle - \langle b, u \rangle$, matrix Q is spd.
 - $\nabla J(u) = Qu - b$, $\|\cdot\|_Q^2 \equiv \langle \cdot, Q \cdot \rangle$.



Gradient descent with exact line search



- Gradient descent with *exact* line search:

$$u^{k+1} = u^k - \tau^k \nabla J(u^k),$$
$$\tau^k = \arg \min_{\tau} J(u^k - \tau \nabla J(u^k)).$$

- Special case: $J(u) = \frac{1}{2} \langle u, Qu \rangle - \langle b, u \rangle$, matrix Q is spd.

- $\nabla J(u) = Qu - b$, $\|\cdot\|_Q^2 \equiv \langle \cdot, Q \cdot \rangle$.

- $\tau^k = \arg \min_{\tau} J(u^k - \tau \nabla J(u^k)) = \frac{\|\nabla J(u^k)\|^2}{\|\nabla J(u^k)\|_Q^2} \Rightarrow$

$$\|u^{k+1} - u^*\|_Q^2 = \left(1 - \frac{\|\nabla J(u^k)\|^4}{\|\nabla J(u^k)\|_Q^2 \|\nabla J(u^k)\|_{Q^{-1}}^2} \right) \|u^k - u^*\|_Q^2$$
$$\leq \left(\frac{\lambda_{\max}(Q) - \lambda_{\min}(Q)}{\lambda_{\max}(Q) + \lambda_{\min}(Q)} \right)^2 \|u^k - u^*\|_Q^2.$$



Backtracking line search

- Sufficient decrease condition (let $c_1 \in (0, 1)$):

$$J(u^k + \tau d^k) \leq J(u^k) + c_1 \tau \langle \nabla J(u^k), d^k \rangle. \quad (\text{A})$$

- Curvature condition (let $c_2 \in (c_1, 1)$):

$$\langle \nabla J(u^k + \tau d^k), d^k \rangle \geq c_2 \langle \nabla J(u^k), d^k \rangle. \quad (\text{C})$$



Backtracking line search

- Sufficient decrease condition (let $c_1 \in (0, 1)$):


$$J(u^k + \tau d^k) \leq J(u^k) + c_1 \tau \langle \nabla J(u^k), d^k \rangle. \quad (\text{A})$$

- Curvature condition (let $c_2 \in (c_1, 1)$):


$$\langle \nabla J(u^k + \tau d^k), d^k \rangle \geq c_2 \langle \nabla J(u^k), d^k \rangle. \quad (\text{C})$$

- (A) \rightsquigarrow **Armijo** line search; (A) & (C) \rightsquigarrow **Wolfe-Powell** I.s.

Armijo I.s.



Wolfe-Powell I.s.



Convergence of backtracking line search

Lemma (feasibility of line search)

Assume that $J : \mathbb{E} \rightarrow \mathbb{R}$ is continuously differentiable, $\langle \nabla J(u^k), d^k \rangle < 0 \forall k$, and $0 < c_1 < c_2 < 1$. Then there exists an open interval in which the step size τ satisfies (A) and (C).

Proof: on board.



Convergence of backtracking line search

Lemma (feasibility of line search)

Assume that $J : \mathbb{E} \rightarrow \mathbb{R}$ is continuously differentiable, $\langle \nabla J(u^k), d^k \rangle < 0 \forall k$, and $0 < c_1 < c_2 < 1$. Then there exists an open interval in which the step size τ satisfies (A) and (C).

Proof: on board.



Theorem (Zoutendijk)

Assume that $J : \mathbb{E} \rightarrow \mathbb{R}$ is cont'ly differentiable, and (A) and (C) are both satisfied with $0 < c_1 < c_2 < 1$ for each k . In addition, J is μ -Lipschitz differentiable on $\{u \in \mathbb{E} : J(u) \leq J(u^0)\}$. Then

$$\sum_{k=0}^{\infty} \frac{|\langle \nabla J(u^k), d^k \rangle|^2}{\|d^k\|^2} < \infty.$$

Proof: on board.

Remark

If $\frac{|\langle \nabla J(u^k), d^k \rangle|}{\|\nabla J(u^k)\| \|d^k\|} \geq \text{constant} > 0$, then $\lim_{k \rightarrow \infty} \|\nabla J(u^k)\| = 0$.

Majorize-minimize method

Majorizing function

A function $\hat{J}(\cdot; u)$ is a **majorant** of J at $u \in \mathbb{E}$ if

$$\begin{cases} \hat{J}(u; u) = J(u), \\ \hat{J}(\cdot; u) \geq J(\cdot). \end{cases}$$



Majorize-minimize method

Majorizing function


A function $\hat{J}(\cdot; u)$ is a **majorant** of J at $u \in \mathbb{E}$ if

$$\begin{cases} \hat{J}(u; u) = J(u), \\ \hat{J}(\cdot; u) \geq J(\cdot). \end{cases}$$

Majorize-minimize (MM) algorithm

Let $\hat{J}(\cdot; u)$ majorize $J \forall u \in \mathbb{E}$. Then the MM iteration reads:

$$u^{k+1} \in \arg \min_u \hat{J}(u; u^k).$$



Remark

- 1 Monotonic decrease of objectives:

$$J(u^{k+1}) \leq \widehat{J}(u^{k+1}; u^k) \leq \widehat{J}(u^k; u^k) = J(u^k).$$

- 2 Efficiency of MM relies on the choice of the majorant $\widehat{J}(\cdot; u)$, i.e., $\widehat{J}(\cdot; u)$ is easy to minimize.
- 3 Common choices of $\widehat{J}(\cdot; u)$ are quadratics.



Remark

- 1 Monotonic decrease of objectives:

$$J(u^{k+1}) \leq \widehat{J}(u^{k+1}; u^k) \leq \widehat{J}(u^k; u^k) = J(u^k).$$

- 2 Efficiency of MM relies on the choice of the majorant $\widehat{J}(\cdot; u)$, i.e., $\widehat{J}(\cdot; u)$ is easy to minimize.
- 3 Common choices of $\widehat{J}(\cdot; u)$ are quadratics.



Gradient descent as MM

- Observe that $u^{k+1} = u^k - \tau \nabla J(u^k)$ iff
$$u^{k+1} = \arg \min_u J(u^k) + \langle \nabla J(u^k), u - u^k \rangle + \frac{1}{2\tau} \|u - u^k\|^2.$$
- When does $J(u^k) + \langle \nabla J(u^k), \cdot - u^k \rangle + \frac{1}{2\tau} \|\cdot - u^k\|^2 \geq J(\cdot)$ hold?

Lemma

Assume that $J : \mathbb{E} \rightarrow \mathbb{R}$ is μ -Lipschitz differentiable. Then
 $\forall u, v \in \mathbb{E}$:

$$|J(v) - J(u) - \langle \nabla J(u), v - u \rangle| \leq \frac{\mu}{2} \|v - u\|^2.$$

Proof: on board.



Lemma

Assume that $J : \mathbb{E} \rightarrow \mathbb{R}$ is μ -Lipschitz differentiable. Then $\forall u, v \in \mathbb{E}$:

$$|J(v) - J(u) - \langle \nabla J(u), v - u \rangle| \leq \frac{\mu}{2} \|v - u\|^2.$$

Proof: on board.

Theorem (convergence of gradient descent)

Assume that $J : \mathbb{E} \rightarrow \mathbb{R}$ is μ -Lipschitz differentiable. Then the gradient descent iteration

$$u^{k+1} = u^k - \tau \nabla J(u^k)$$

with $\tau \in (0, 1/\mu]$ yields $\lim_{k \rightarrow \infty} \nabla J(u^k) = 0$.

Proof: on board.



Gradient descent as MM

Lemma

Assume that $J : \mathbb{E} \rightarrow \mathbb{R}$ is μ -Lipschitz differentiable. Then $\forall u, v \in \mathbb{E}$:

$$|J(v) - J(u) - \langle \nabla J(u), v - u \rangle| \leq \frac{\mu}{2} \|v - u\|^2.$$

Proof: on board.



Theorem (convergence of gradient descent)

Assume that $J : \mathbb{E} \rightarrow \mathbb{R}$ is μ -Lipschitz differentiable. Then the gradient descent iteration

$$u^{k+1} = u^k - \tau \nabla J(u^k)$$

with $\tau \in (0, 1/\mu]$ yields $\lim_{k \rightarrow \infty} \nabla J(u^k) = 0$.

Proof: on board.

Recipe of (global) convergence

By solving the surrogate problem in MM, we achieve: (1) sufficient decrease in the objective; (2) inexact optimality condition matches the exact OC in the limit.