



Multiple View Geometry: Solution Sheet 6

Prof. Dr. Florian Bernard, Lukas Köstler, Mohammed Brahim
Computer Vision Group, TU Munich

<http://vision.in.tum.de/teaching/ss2020/mvg2020>

Exercise: June 4th, 2020

Part I: Theory

1. (a) E is essential matrix $\Rightarrow \Sigma = \text{diag}\{\sigma, \sigma, 0\}$:

$$R_z(\pm\frac{\pi}{2})\Sigma = \begin{pmatrix} 0 & \mp 1 & 0 \\ \pm 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} \sigma & 0 & 0 \\ 0 & \sigma & 0 \\ 0 & 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & \mp\sigma & 0 \\ \pm\sigma & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} = -(R_z(\pm\frac{\pi}{2})\Sigma)^\top$$

$$\begin{aligned} -\hat{T}^\top &= -(UR_z\Sigma U^\top)^\top \\ &= U(-R_z\Sigma)^\top U^\top \\ &= UR_z\Sigma U^\top \\ &= \hat{T} \end{aligned}$$

- (b) Since U, V are orthogonal with determinant 1 (see lecture), they are rotation matrices. Since $\text{SO}(3)$ is a group and thus closed under multiplication, $R \in \text{SO}(3)$.

Alternative longer proof:

- i. U, V are orthogonal matrices $\Rightarrow U^\top U = \mathbb{1}$ and $VV^\top = \mathbb{1}$
 R_z is a rotation matrix $\Rightarrow R_z R_z^\top = \mathbb{1}$

$$\begin{aligned} R^\top R &= (UR_z^\top V^\top)^\top (UR_z^\top V^\top) \\ &= VR_z U^\top UR_z^\top V^\top \\ &= VR_z R_z^\top V^\top \\ &= VV^\top \\ &= \mathbb{1} \end{aligned}$$

- ii. U and V are special orthogonal matrices with $\det(U) = \det(V^\top) = 1$ (Slide 9, Chapter 5).

$$\det(R) = \det(UR_z^\top V^\top) = \underbrace{\det(U)}_1 \cdot \underbrace{\det(R_z^\top)}_1 \cdot \underbrace{\det(V^\top)}_1 = 1$$

2. (a) $H = R + Tu^\top \Leftrightarrow R = H - Tu^\top$.

$$\begin{aligned} E &= \hat{T}R \\ &= \hat{T}(H - Tu^\top) \\ &= \hat{T}H - \underbrace{\hat{T}T}_{=T \times T=0} u^\top \\ &= \hat{T}H \end{aligned}$$

(b)

$$\begin{aligned}
H^\top E + E^\top H &= H^\top (\hat{T}H) + (\hat{T}H)^\top H \\
&= H^\top (\hat{T}H) + H^\top \hat{T}^\top H \\
&= H^\top \hat{T}H - H^\top \hat{T}H \quad (\text{because } \hat{T} \text{ is skew-symmetric, i.e. } \hat{T}^\top = -\hat{T}) \\
&= 0
\end{aligned}$$

3. The notations below are as in Slide 6, Chapter 5. Note that the following slides deal with projected points in the normalized plane ($Z = 1$), whereas here we assume pixel coordinates. The case of normalized coordinates is then just a special case with $K = \mathbb{1}$.

Rotation R and translation T are defined such that

$$g_{21} = \begin{bmatrix} R & T \\ 0 & 1 \end{bmatrix}$$

transforms a point from coordinate system 1 (CS1) to coordinate system 2 (CS2). This means that the inverse transformation (converting points from CS2 to CS1) is given by

$$g_{12} = g_{21}^{-1} = \begin{bmatrix} R^\top & -R^\top T \\ 0 & 1 \end{bmatrix}.$$

o_1 seen in CS1: $[0 \ 0 \ 0 \ 1]^\top$ (homogeneous coordinates)

o_1 seen in CS2: $g_{21} [0 \ 0 \ 0 \ 1]^\top = \begin{bmatrix} T \\ 1 \end{bmatrix}$

e_2 are the pixel coordinates of o_1 projected into image 2:

$$\lambda_1 e_2 = K \Pi_0 \begin{bmatrix} T \\ 1 \end{bmatrix} = KT$$

o_2 seen in CS2: $[0 \ 0 \ 0 \ 1]^\top$

o_2 seen in CS1: $g_{12} [0 \ 0 \ 0 \ 1]^\top = \begin{bmatrix} -R^\top T \\ 1 \end{bmatrix}$

e_1 are the pixel coordinates of o_2 projected into image 1:

$$\lambda_2 e_1 = K \Pi_0 \begin{bmatrix} -R^\top T \\ 1 \end{bmatrix} = -KR^\top T$$

$$\begin{aligned}
Fe_1 &= \underbrace{(K^{-\top} \hat{T} R K^{-1})}_F \underbrace{\left(-\frac{1}{\lambda_2} K R^\top T\right)}_{e_1} \\
&= -\frac{1}{\lambda_2} K^{-\top} \hat{T} R \underbrace{K^{-1} K}_\mathbb{1} R^\top T \\
&= -\frac{1}{\lambda_2} K^{-\top} \hat{T} \underbrace{R R^\top}_\mathbb{1} T \\
&= -\frac{1}{\lambda_2} K^{-\top} \underbrace{\hat{T} T}_{=T \times T=0} \\
&= 0
\end{aligned}$$

$$\begin{aligned}
e_2^\top F &= \underbrace{\left(\frac{1}{\lambda_1}KT\right)^\top}_{e_2} \underbrace{\left(K^{-\top}\hat{T}RK^{-1}\right)}_F \\
&= \frac{1}{\lambda_1}T^\top \underbrace{K^\top K^{-\top}}_{\mathbb{1}} \hat{T}RK^{-1} \\
&= \frac{1}{\lambda_1}T^\top \hat{T}RK^{-1} \\
&= \frac{1}{\lambda_1}(\hat{T}^\top T)^\top RK^{-1} \\
&= \frac{1}{\lambda_1}(-\hat{T}T)^\top RK^{-1} \\
&= -\frac{1}{\lambda_1}(T \times T)^\top RK^{-1} \\
&= -\frac{1}{\lambda_1}0RK^{-1} \\
&= 0
\end{aligned}$$

Part II: Practical Exercises

Remarks

1. Are there two or four possible solutions for R and T ?

The answer is four. You may find it confusing since in Slide 9 you are told to have two sets of R and T , while in Slide 14 it says there are four possible solutions. Recall that the essential matrix E is calculated by solving the equation $\chi E^s = 0$ (at the bottom of Slide 11). Ideally the E^s you get should lie in the nullspace of χ . However, since there are always errors in the point pairs you've chosen to make χ , in practice it is very difficult to get an E^s that makes χE^s exactly 0. Instead we use the SVD of χ to get the E^s which minimizes $\|\chi E^s\|$. In other words, the E^s we get will give us $\chi E^s = \sigma$ with σ being some very small vector. Now think about this: what will happen if we turn the sign of the E^s ? We will get $\chi(-E^s) = -\sigma$ which still gives us the smallest $\|\chi E^s\|$ ($\|\chi E^s\| = \|\sigma\| = \|-\sigma\| = \|\chi(-E^s)\|$).

Now we know that solving $\chi E^s = 0$ for E^s will always give us two possible solutions E and $-E$. From each of them you can get two possible sets of R and T using the equations in Slide 9. Altogether we get four possible solutions. In practice we do the calculation in a little different way. We usually calculate according to Slide 14 to get four solutions out of E . The two extra solutions we get are nothing but the ones we should have got from $-E$.

2. How to get the correct R and T from the possible solutions?

The criterion you should use to rule out the incorrect solutions is that all the reconstructed 3D points should have positive depth seen from **both of the camera coordinate systems**. In other words, both λ_1^j and λ_2^j in Slide 17 need to be positive.