



## Multiple View Geometry: Solution Sheet 2

Prof. Dr. Daniel Cremers,

Shenhan Qian, Simon Weber, Anna Ribic, and Tarun Yenamandra

Computer Vision Group, TU Munich

Wednesdays 16:15–18:15 at Hörsaal 2, "Interims I"  
(5620.01.102), and on RBG Live

Exercise: May 14th, 2025

1. Rigid body motion requires to preserve 1) the norm and 2) the cross product. We first show the norm preservation, given a vector  $v \in \mathbb{R}^3$  and a rotation matrix  $R \in \mathbb{R}^{3 \times 3}$ :

$$\|Rv\|^2 = (Rv)^\top Rv = v^\top R^\top Rv = v^\top v = \|v\|^2$$

Next, we show the cross product preservation. Given two vectors  $a$  and  $b$  and assume we only rotate around x-axis with angle  $\theta$ . Therefore, the rotation matrix is:

$$R_x(\theta) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos(\theta) & -\sin(\theta) \\ 0 & \sin(\theta) & \cos(\theta) \end{pmatrix}$$

Rotating the cross product of  $a$  and  $b$  with  $R_x(\theta)$ , we have:

$$\begin{aligned} a \times b &= \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix} \times \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix} = \begin{pmatrix} a_2 b_3 - a_3 b_2 \\ a_3 b_1 - a_1 b_3 \\ a_1 b_2 - a_2 b_1 \end{pmatrix} \\ \Rightarrow R_x(\theta)(a \times b) &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos(\theta) & -\sin(\theta) \\ 0 & \sin(\theta) & \cos(\theta) \end{pmatrix} \begin{pmatrix} a_2 b_3 - a_3 b_2 \\ a_3 b_1 - a_1 b_3 \\ a_1 b_2 - a_2 b_1 \end{pmatrix} \\ &= \begin{pmatrix} a_2 b_3 - a_3 b_2 \\ \cos(\theta)(a_3 b_1 - a_1 b_3) - \sin(\theta)(a_1 b_2 - a_2 b_1) \\ \sin(\theta)(a_3 b_1 - a_1 b_3) + \cos(\theta)(a_1 b_2 - a_2 b_1) \end{pmatrix} \end{aligned}$$

Now, we first rotate  $a$  and  $b$  with  $R_x(\theta)$  and then calculate the cross product:

$$\begin{aligned} (R_x(\theta)a) \times (R_x(\theta)b) &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos(\theta) & -\sin(\theta) \\ 0 & \sin(\theta) & \cos(\theta) \end{pmatrix} \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix} \times \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos(\theta) & -\sin(\theta) \\ 0 & \sin(\theta) & \cos(\theta) \end{pmatrix} \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix} \\ &= \begin{pmatrix} a_1 \\ \cos(\theta)a_2 - \sin(\theta)a_3 \\ \sin(\theta)a_2 + \cos(\theta)a_3 \end{pmatrix} \times \begin{pmatrix} b_1 \\ \cos(\theta)b_2 - \sin(\theta)b_3 \\ \sin(\theta)b_2 + \cos(\theta)b_3 \end{pmatrix} \end{aligned}$$

Following the calculation of the cross product, we can see that the two results are the same. Therefore, the cross product is preserved under rotation around the x-axis. The same can be shown for the y and z axis. The corresponding rotation matrices are:

$$R_y(\theta) = \begin{pmatrix} \cos(\theta) & 0 & \sin(\theta) \\ 0 & 1 & 0 \\ -\sin(\theta) & 0 & \cos(\theta) \end{pmatrix} \quad R_z(\theta) = \begin{pmatrix} \cos(\theta) & -\sin(\theta) & 0 \\ \sin(\theta) & \cos(\theta) & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$2. \lambda_a = \frac{(\lambda_a v_a)^\top v_b}{\langle v_a, v_b \rangle} = \frac{v_a^\top A^\top v_b}{\langle v_a, v_b \rangle} = \frac{v_a^\top A v_b}{\langle v_a, v_b \rangle} = \frac{v_a^\top (\lambda_b v_b)}{\langle v_a, v_b \rangle} = \lambda_b$$

3. We assume  $u \neq -v$ . Otherwise, we only get  $w = 0$ . We first compute the inner product between  $u$  and  $v$ :

$$\langle u, v \rangle = \|u\| \|v\| \cos(\theta) = 1 \cdot 1 \cdot \cos(\theta) = \cos(\theta)$$

Next, we compute the inner product between  $w$  and  $u$ :

$$\langle w, u \rangle = \langle u + v, u \rangle = \langle u, u \rangle + \langle v, u \rangle = 1 + \cos(\theta)$$

Alternatively, we can compute the inner product between  $w$  and  $u$  as follows:

$$\langle w, u \rangle = \|w\| \|u\| \cos(\alpha) = \|u + v\| \cos(\alpha)$$

Now, we compute  $\|u + v\|$ :

$$\|u + v\| = \sqrt{\langle u + v, u + v \rangle} = \sqrt{\langle u, u \rangle + 2\langle u, v \rangle + \langle v, v \rangle} = \sqrt{1 + 2\cos(\theta) + 1} = \sqrt{2(1 + \cos(\theta))}$$

Finally, we compute  $\cos(\alpha)$ :

$$\begin{aligned} \|u + v\| \cos(\alpha) &= 1 + \cos(\theta) \\ \Rightarrow \sqrt{2(1 + \cos(\theta))} \cos(\alpha) &= 1 + \cos(\theta) \\ \Rightarrow \cos(\alpha) &= \frac{1 + \cos(\theta)}{\sqrt{2(1 + \cos(\theta))}}. \end{aligned}$$

Because  $\cos^2(\theta/2) = \frac{1 + \cos(\theta)}{2}$ , we have shown  $\cos(\alpha) = \cos(\theta/2)$ .

4. Let  $V$  be the orthonormal matrix (i.e.  $V^\top = V^{-1}$ ) given by the eigenvectors, and  $\Sigma$  the diagonal matrix containing the eigenvalues:

$$V = \begin{pmatrix} | & & | \\ v_1 & \cdots & v_n \\ | & & | \end{pmatrix} \quad \text{and} \quad \Sigma = \begin{pmatrix} \lambda_1 & 0 & \ddots \\ 0 & \ddots & 0 \\ \ddots & 0 & \lambda_n \end{pmatrix}.$$

As  $V$  is a basis, we can express  $x$  as a linear combination of the eigenvectors  $x = V\alpha$ , for some  $\alpha \in \mathbb{R}^n$ . For  $\|x\| = 1$  we have  $\sum_i \alpha_i^2 = \alpha^\top \alpha = x^\top V V^\top x = x^\top x = 1$ . This gives

$$\begin{aligned} x^\top A x &= x^\top V \Sigma V^{-1} x \\ &= \alpha^\top V^\top V \Sigma V^\top V \alpha \\ &= \alpha^\top \Sigma \alpha = \sum_i \alpha_i^2 \lambda_i \end{aligned}$$

Considering  $\sum_i \alpha_i^2 = 1$ , we can conclude that this expression is minimized iff only the  $\alpha_i$  corresponding to the smallest eigenvalue(s) are non-zero. If  $\lambda_{n-1} \geq \lambda_n$ , there exist only two solutions ( $\alpha_n = \pm 1$ ), otherwise infinitely many.

For maximisation, only the the  $\alpha_i$  corresponding to the largest eigenvalue(s) can be non-zero.

5. We show that:  $x \in \text{kernel}(A) \Leftrightarrow x \in \text{kernel}(A^\top A)$ .

" $\Rightarrow$ ": Let  $x \in \text{kernel}(A)$

$$A^\top \underbrace{Ax}_{=0} = A^\top 0 = 0 \Rightarrow x \in \text{kernel}(A^\top A)$$

" $\Leftarrow$ ": Let  $x \in \text{kernel}(A^\top A)$

$$0 = x^\top \underbrace{A^\top Ax}_{=0} = \langle Ax, Ax \rangle = \|Ax\|^2 \Rightarrow Ax = 0 \Rightarrow x \in \text{kernel}(A)$$

## 6. Singular Value Decomposition (SVD)

*Note:* There exist multiple slightly different definitions of the SVD. Depending on the convention used, we might have  $S \in \mathbb{R}^{m \times n}$ ,  $S \in \mathbb{R}^{n \times n}$ , or  $S \in \mathbb{R}^{p \times p}$  where  $p = \text{rank}(A)$ . In the lecture the last option was presented. In the following, we present the results for the first option, since that is the one that `numpy.linalg.svd` function returns by default.

(a)  $A \in \mathbb{R}^{m \times n}$ ,  $U \in \mathbb{R}^{m \times m}$ ,  $S \in \mathbb{R}^{m \times n}$ ,  $V \in \mathbb{R}^{n \times n}$

(b) Similarities and differences between SVD and EVD:

i. Both are matrix diagonalization techniques.

ii. The SVD can be applied to matrices  $A \in \mathbb{R}^{m \times n}$  with  $m \neq n$ , whereas the EVD is only applicable to quadratic matrices ( $A \in \mathbb{R}^{m \times n}$  with  $m = n$ ).

(c) Relationship between  $U$ ,  $S$ ,  $V$  and the eigenvalues and eigenvectors of  $A^\top A$  and  $AA^\top$ :

i.  $A^\top A$ : The columns of  $V$  are eigenvectors; the squares of the diagonal elements of  $S$  are eigenvalues.

ii.  $AA^\top$ : The columns of  $U$  are eigenvectors; the squares of the diagonal elements of  $S$  are eigenvalues (possibly filled up with zeros).

(d) Entries in  $S$ :

i.  $S$  is a diagonal matrix. The elements along the diagonal are the *singular values* of  $A$ .

ii. The number of non-zero singular values gives us the *rank* of the matrix  $A$ .