Chapter 2 **Diffusion Filtering**

Computer Vision I: Variational Methods

Winter 2017/18

Dr. Yvain Quéau Chair for Computer Vision and Pattern Recognition Departments of Informatics & Mathematics Technical University of Munich Diffusion Filtering

Dr Yvain Qué Au



Nonlinear Filtering

Partial Differential Equations

Diffusion

Overview

Diffusion Filtering

Dr. Yvain QuÉAU



1 Nonlinear Filtering

Nonlinear Filtering Partial Differential

Equations

Diffusion

Nonlinear & Anisotropic Diffusion

2 Partial Differential Equations

Diffusion

Overview

Diffusion Filtering

Dr. Yvain QuÉAU



Nonlinear Filtering

Nonlinear Filtering

Partial Differential Equations

Diffusion

Nonlinear & Anisotropic Diffusion

2 Partial Differential Equations

Nonlinear Filtering

• The convolution of an input image f(x) with a kernel G(x):

$$g(x) = (G * f)(x) = \int G(x')f(x - x')dx'$$

is a classical example of a linear filter.

- Convolutions can be efficiently implemented in frequency space because in frequency space the convolution corresponds to a simple (frequency-wise) product and because the Fast Fourier transform allows a quick conversion to and from frequency space.
- In practice, however, linear filters are often suboptimal. In smoothing/denoising, for example, the Gaussian smoothing removes both noise and signal semantically relevant structures tend to disappear along with the noise. Instead, one would like to remove noise in an adaptive manner such that semantically important structures remain unaffected. In principle this could be done with a Gaussian smoothing where the filter width σ is adapted to the local structure (larger in noise areas, smaller at important edges).

Diffusion Filtering

Dr. Yvain QuÉAU



Nonlinear Filtering

Partial Differential Equations

Diffusion

Image Filtering by Diffusion

Formally this would amount to the following:

$$g(x) = \int G_{\sigma(f,x)}(x')f(x-x')dx',$$

where now the width σ of the convolution kernel G depends on the brightness values in a local neighborhood.

- It turns out that there exist other more elegant solutions to model such adaptive denoising processes by means of Diffusion filtering.
- The key observation is that image smoothing can be modeled with a diffusion process. In this process, the local brightness diffuses to neighboring pixels due to differences in the local concentration of grayvalue.
- Mathematically diffusion processes are represented by partial differential equations (PDEs).

Diffusion Filtering

Dr Yvain Qué Au



Ionlinear Filtering

Partial Differential Equations

Diffusion

Overview

Diffusion Filtering

Dr. Yvain QuÉAU



Nonlinear Filtering

Nonlinear Filtering

Partial Differential Equations

Diffusion

Nonlinear & Anisotropic Diffusion

- 2 Partial Differential Equations

Review: Partial Differential Equations

 A partial differential equation (PDE) is an equation containing the partial derivatives of a function of several variables

Example — the wave equation:

$$\frac{\partial^2 \psi(\mathbf{x},t)}{\partial t^2} = c^2 \Delta \psi(\mathbf{x},t)$$

 For functions of a single variable we have the special case of ordinary differential equations (ODEs) (gewöhnliche Differentialgleichungen).

Example — the pendulum:

$$m\frac{d^2x(t)}{dt^2} + \gamma\frac{dx(t)}{dt} + kx(t) = 0$$

 Many natural phenomena can be modeled by partial differential equations. In most cases, one can derive the respective equation from a few basic principles. A solution of a differential equation is a function for which the differential equation is true.

Diffusion Filtering

Dr Yvain Qué Au



Nonlinear Filtering

Diffusion

Analytical Solutions

- A few PDEs can be solved analytically, i.e. the solution can be written in closed form.
- Example The wave equation (in 1D):

$$\frac{\partial^2 \psi(x,t)}{\partial t^2} = c^2 \frac{\partial^2 \psi(x,t)}{\partial x^2}$$

has the (not necessarily unique) solution: $\psi(x, t) = \sin(x - ct)$

- If solutions are not unique one can impose additional assumptions boundary conditions or initial conditions, for example $\psi(x,0) = \psi_0(x)$.
- Example The harmonic oscillator (without friction):

$$m\frac{d^2x(t)}{dt^2}+kx(t)=0$$

has the (generally not unique) solution:

$$x(t) = \sin(\omega t)$$
, with $\omega = \sqrt{k/m}$.

Diffusion Filtering

Dr. Yvain QuÉAU



Nonlinear Filtering

Partial Differential Equations

Diffusion

Non-Analytical Solutions - The Finite Differences Method

Example: discretization of the 2D wave equation

$$\frac{\partial^2 u(x,y,t)}{\partial t^2} = c^2 \Delta u(x,y,t)$$

Time discretization (backward, order 2):

$$\begin{array}{l} \frac{\partial^2 u(\cdot,\cdot,t)}{\partial t^2} \approx u^{(t)} - 2u^{(t-1)} + u^{(t-2)} \\ \Rightarrow u^{(t)} - 2u^{(t-1)} + u^{(t-2)} = c^2 \Delta u^{(t)} \\ \Rightarrow \left(\mathrm{id} - c^2 \Delta\right) u^{(t)} = 2u^{(t-1)} - u^{(t-2)} \\ (\mathrm{assume} \ u^{(-1)} \ \mathrm{and} \ u^{(0)} \ \mathrm{are} \ \mathrm{known} \ \mathrm{-initial} \ \mathrm{condition}) \end{array}$$

• Space discretization (central, order 2): $\Delta u(x,y,\cdot) \approx u(x+1,y) + u(x-1,y) + u(x,y+1) + u(x,y-1) - 4u(x,y)$ (assume, e.g., Dirichlet boundary conditions)

 \Rightarrow Linear system $A^{(t)}u^{(t)}(:) = b^{(t)}$ to solve to obtain u^t

Diffusion Filtering

Dr. Yvain QuÉAU



Nonlinear Filtering

Partial Differential Equations

Diffusion

Example: numerical simulation of the 2D wave equation

Diffusion Filtering

Dr. Yvain QuÉAU



Nonlinear Filtering

Partial Differential Equations

Diffusion

Overview

Diffusion Filtering

Dr. Yvain QuÉAU



Nonlinear Filtering

Nonlinear Filtering

Partial Differential Equations

2 Partial Differential Equations

Nonlinear & Anisotropic Diffusion

Diffusion

The Diffusion Equation

- Diffusion is a physical process which aims at minimizing differences in the spatial concentration u(x, t) of a substance.
- This process can be described by two basic equations:
 - <u>Fick's law</u> states that concentration differences induce a flow j of the substance in direction of the negative concentration gradient:

$$j = -g\nabla u$$

The diffusivity g describes the speed of the diffusion process.

The continuity equation

$$\partial_t u = -\operatorname{div} i$$

where div $j \equiv \nabla^{\top} j \equiv \partial_x j_1 + \partial_y j_2$ is called the divergence of the vector j.

Inserting one into the other leads to the diffusion equation:

$$\partial_t u = \operatorname{div}\left(g\cdot\nabla u\right)$$

Diffusion Filtering

Dr. Yvain QuÉAU



Nonlinear Filtering

Partial Differential Equations

ffusion

Solution of the Linear Diffusion Equation

The one-dimensional linear diffusion equation (g = 1)

$$\partial_t u = \partial_x^2 u.$$

with initial condition

$$u(x,t=0)=f(x)$$

has the unique solution:

$$u(x,t)=(G_{\sqrt{2t}}*f)(x)=\int_{-\infty}^{\infty}G_{\sqrt{2t}}(x-x')f(x')dx',$$

where

$$G_{\sigma}(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{x^2}{2\sigma^2}},$$

is a Gaussian kernel of width $\sigma = \sqrt{2t}$.

Diffusion Filtering

Dr. Yvain Quéau



Nonlinear Filtering

Partial Differential Equations

iffusion



Nonlinear Filtering

Partial Differential Equations

Diffusion

- The above result implies that smoothing of an image $f: \Omega \subset \mathbb{R}^2 \to \mathbb{R}$ with Gaussian kernels of increasing width σ can be realized through a diffusion process of the form
 - $\begin{cases} \partial_t u(x,t) = \Delta u \\ u(x,0) = f(x) \quad \forall x \in \Omega \\ \partial_n u|_{\partial\Omega} = \langle \nabla u, n \rangle|_{\partial\Omega} = 0 \end{cases}$
- The latter boundary condition states that the derivative of the brightness function u along the normal n at the image boundary $\partial\Omega$ must vanish. This assures that no brightness will leave or enter the image, i.e. the average brightness will be preserved.
- With increasing time t the solution u(x,t) of this process will correspond to increasingly smoothed versions of the original image f(x).

Equations

Nonlinear & Anisotropic Diffusion

-

Discretization of the 2D diffusion equation

 $\frac{\partial u(x,y,t)}{\partial t} = \text{div } (g \cdot \nabla u(x,y,t))$

Time discretization (backward, order 1):

$$\begin{array}{l} \frac{\partial u(\cdot,\cdot,t)}{\partial t} \approx u^{(t)} - u^{(t-1)} \\ \Rightarrow \left(\mathrm{id} - \mathrm{div} \left(g \cdot \nabla \right) \right) u^{(t)} = u^{(t-1)} \\ (\mathrm{assume} \ u^{(0)} \ \mathrm{known} \ - \ \mathrm{initial} \ \mathrm{condition}) \end{array}$$

- Space discretization (order 1 forward for the gradient, backward for the divergence):
 - \Rightarrow Linear system $A^{(t)}u^{(t)}(:) = b^{(t)}$ to solve to obtain u^t

Example: numerical simulation of the 2D diffusion

With uniform diffusivity ($g \equiv 1$)

Diffusion Filtering

Dr. Yvain QuÉAU



Nonlinear Filtering
Partial Differential

Partial Differential Equations

Diffusion

Example: numerical simulation of the 2D diffusion

With non-uniform diffusivity ($g \equiv 1$ except on the "walls")

Diffusion Filtering

Dr. Yvain QuÉAU



Nonlinear Filtering

Partial Differential Equations

Diffusion

Overview

Diffusion Filtering

Dr. Yvain QuÉAU



Nonlinear Filtering

Partial Differential Equations

Diffusion

Nonlinear Filtering

2 Partial Differential Equations

Nonlinear and Anisotropic Diffusion

General diffusion equation:

$$\partial_t u = \operatorname{div}(g \nabla u)$$

- For g=1 (or $g=\text{const.} \in \mathbb{R}$) the diffusion process is called linear, isotropic and homogeneous.
- If the diffusivity g is space-dependent, i.e. g = g(x), the process is called an inhomogeneous diffusion.
- If the diffusivity depends on u, i.e. g = g(u), then it is called a nonlinear diffusion because then the equation is no longer linear in u.
- If the diffusivity g is matrix-valued then the process is called an anisotropic diffusion. A matrix-valued diffusivity leads to processes where the diffusion is different in different directions.
- Note: In the literature this terminology is not used consistently.

Diffusion Filtering

Dr. Yvain QuÉAU



Nonlinear Filtering
Partial Differential

Equations Diffusion

Edge-preserving Diffusion

 Idea: Less diffusion (smoothing) in locations of strong edge information.

- Gradient norm $|\nabla u| = \sqrt{u_{x}^{2} + u_{y}^{2}}$ serves as edge indicator
- Diffusivity should decrease with increasing |∇u|. For example (Perona & Malik, Scale Space and Edge Detection using Anisotropic Diffusion, PAMI 1990):

$$g(|\nabla u|) = \exp\left\{-|\nabla u|^2/\lambda^2\right\} \text{ or } g(|\nabla u|) = \frac{1}{1+|\nabla u|^2/\lambda^2}$$

- $\lambda>0$ is called a contrast parameter. Areas where $|\nabla u|\gg\lambda$ will not be affected much by the diffusion process.
- The Perona-Malik model had a huge impact in image processing because it allowed a better edge detection than classical edge detectors (such as the Canny edge detector).

Diffusion Filtering

Dr. Yvain QuÉAU



Nonlinear Filtering

Partial Differential Equations

Diffusion

With uniform diffusivity ($g \equiv 1$)

Diffusion Filtering

Dr. Yvain QuÉAU



Nonlinear Filtering

Partial Differential Equations

Diffusion

With Perona-Malik's "anisotropic" diffusivity $(g = \exp\{-|\nabla u|^2/\lambda^2\}), \lambda = 5$

Diffusion Filtering

Dr. Yvain QuÉAU



Nonlinear Filtering

Partial Differential Equations

Diffusion

With Perona-Malik's "anisotropic" diffusivity ($g = \frac{1}{1 + |\nabla u|^2/\lambda^2}$), $\lambda = 5$

Diffusion Filtering

Dr. Yvain QuÉAU



Nonlinear Filtering

Partial Differential Equations

Diffusion

With minimal surface diffusivity (
$$g=rac{1}{\sqrt{1+|
abla u|^2/\lambda^2}}$$
), $\lambda=1$

Diffusion Filtering

Dr. Yvain QuÉAU



Nonlinear Filtering

Partial Differential Equations

Diffusion

Appropriately Setting the Contrast Parameter λ is Difficult

Example with $g = \frac{1}{1 + |\nabla u|^2/\lambda^2}$

Diffusion Filtering

Dr. Yvain QuÉAU



Nonlinear Filtering

Partial Differential Equations

Diffusion

Modifying the Anisotropic Diffusion PDE

Example, prevent diffusion to goo "too far" from u_0 :

$$\partial_t u = \operatorname{div}\left(g\nabla u\right) - \mu\left(u - u_0\right)$$

Diffusion Filtering

Dr. Yvain QuÉAU



Nonlinear Filtering

Partial Differential Equations

Diffusion



Nonlinear Filtering

Partial Differential Equations Diffusion

Time discretization of:

$$\partial_t u = \operatorname{div}(g\nabla u)$$

Explicit time discretization (Stepsize must be small!):

$$\frac{u^{(t+1)}-u^{(t)}}{\Delta t}=\operatorname{div}\left(g\nabla u^{(t)}\right)\Rightarrow u^{(t+1)}=u^{(t)}+\Delta t\operatorname{div}\left(g\nabla u^{(t)}\right)$$

 Implicit time discretization (allow large stepsize, but requires a linear system to be solved):

$$\frac{u^{(t+1)}-u^{(t)}}{\Delta t}=\operatorname{div}\left(g\nabla u^{(t+1)}\right)\Rightarrow\left(\operatorname{id}-\Delta t\operatorname{div}\left(g\nabla\right)\right)u^{(t+1)}=u^{(t)}$$

Explicit VS Implicit Time Discretization

Example of explicit time discretization, with $g=\frac{1}{1+|\nabla u|^2/\lambda^2}$, $\lambda=5$, and stepsize $\Delta t=1$:

 $+|\nabla u|^2/\lambda^2$,

Nonlinear Filtering

Diffusion Filtering

Dr. Yvain QuÉAU

Partial Differential Equations

Diffusion

$$\partial_t u = \partial_x \left(g(|\nabla u|) \partial_x u \right) + \partial_y \left(g(|\nabla u|) \partial_y u \right)$$

Discretize the operators as:

$$\partial_t u pprox rac{u_{ij}^{t+1} - u_{ij}^t}{ au}$$

and

$$\partial_{x}(g\partial_{x}u) \approx \left((g\partial_{x}u)_{i+1/2,j}^{t} - (g\partial_{x}u)_{i-1/2,j}^{t} \right) \\ \approx \left(g_{i+1/2,j}^{t}(u_{i+1,j}^{t} - u_{ij}^{t}) - g_{i-1/2,j}^{t}(u_{ij}^{t} - u_{i-1,j}^{t}) \right)$$

where $g_{i+1/2,j} = \sqrt{g_{i+1,j}g_{ij}}$ assures that no diffusion takes place as soon as g is zero at one of the two pixels.

- Insert, solve for u_{ii}^{t+1} and iterate in t.
- Source: J. Weickert, *Anisotropic Diffusion in Image Processing*.

Diffusion Filtering

Dr. Yvain QuÉAU



Nonlinear Filtering

Partial Differential Equations

Diffusion

Nonlinear Diffusion a la Weickert



Dr. Yvain QuÉAU



Partial Differential Equations Diffusion

Nonlinear Filtering









Lena original





Author: D. Cremers