Image Segmentation II Variational Approache

Dr. Yvain QuÉAU



Variational Image Segmentation

Snakes

The Mumford-Shah Model

Euler-Lagrange Equations

Gradient Descent Implementations

Dr. Yvain Quéau Chair for Computer Vision and Pattern Recognition Departments of Informatics & Mathematics Technical University of Munich

Computer Vision I: Variational Methods

Winter 2017/18

The Mumford-Shah Model

Euler-Lagrange Equations

Gradient Descent Implementations

1 Variational Image Segmentation

Snakes

3 The Mumford-Shah Model

4 Euler-Lagrange Equations

6 Gradient Descent

6 Implementations

Euler-Lagrange Equations

Gradient Descent

Implementations

1 Variational Image Segmentation

The Mumford-Shah Model

Euler-Lagrange Equations

We already studied a number of segmentation algorithms. They were based on two complementary concepts:

- Detecting discontinuities of the brighness function, or
- Grouping pixels of similar brightness (color, texture, etc.)

Most of the approaches discussed so far lack a clear optimization criterion: Edge regions are heuristically fused to connected lines (Perkins, Canny), or pixels are iteratively merged to regions (region merging, region growing).

Toward the end of the 1980s, the first variational formulations for image segmentation emerged, in particular:

- the Snakes (Kass, Witkin, Terzopoulos, "Snakes: Active contour models", Int. J. of Comp. Vision '88),
- the Mumford-Shah Functional (Mumford, Shah, "Optimal approximations by piecewise smooth functions and associated variational problems", J. Appl. Math. '89).

Image Segmentation II Variational Approache

Dr. Yvain QuÉAU



The Mumford-Shah

Euler-Lagrange Equations

Euler-Lagrange Equations

Gradient Descent

Implementations

Variational Image Segmentation

Snakes

The Mumford-Shah Model

Euler-Lagrange Equations

$$E(C) = E_{ext}(C) + E_{int}(C)$$

with an external energy

$$E_{\mathrm{ext}}(C) = -\int\limits_0^1 |\nabla I(C(s))|^2 ds$$

and an internal energy

$$E_{int}(C) = \int_{0}^{1} \left\{ \frac{\alpha}{2} \left| C_{s}(s) \right|^{2} + \frac{\beta}{2} \left| C_{ss}(s) \right|^{2} \right\} ds$$

Here, $I:\Omega\subset\mathbb{R}^2\to\mathbb{R}$ denotes the input image, and $C: [0,1] \to \Omega$ denotes a parametric curve. C_s and C_{ss} denote the first and second derivative of the curve C with respect to its parameter s.

Image Segmentation II Variational Approaches

Dr. Yvain QuÉAU



Variational Image Seamentation

The Mumford-Shah

Euler-Lagrange Equations Gradient Descent

Implementations

$$E_{ext}(C) = -\int_{-\infty}^{1} |\nabla I(C)|^2 ds$$

measures for a given curve C how well it coincides with the maxima of the brightness gradient $|\nabla I|$.

Thus rather than first searching for these maxima and then grouping them to a curve one defines a cost function which measures the "edge strength" along any conceivable curve.

Subsequently, the optimal curve $\hat{\textit{C}}$ is determined by minimizing the total energy:

$$\hat{C} = \arg\min_{C} E(C)$$

Gradient descent on this energy induces an evolution of the curve toward locations of large image gradient.

Image Segmentation II: Variational Approaches

Dr. Yvain QuÉAU



Variational Image Segmentation

kes

The Mumford-Shah Model

Euler-Lagrange Equations Gradient Descent

Implementations

Snakes

The Mumford-Shah Model

Euler-Lagrange Equations Gradient Descent

Implementations

The internal energy is a regularizer which induces some smoothness on the computed curves:

$$E_{int}(C) = \int\limits_0^1 \left\{ rac{lpha}{2} ig| C_{s}(s) ig|^2 + rac{eta}{2} ig| C_{ss}(s) ig|^2
ight\} ds$$

It consists of two components, weighted by parameters $\alpha \geq 0$ and $\beta \geq 0$, which penalize the elastic length and the stiffness of the curve.

Minimizing the total energy

$$E(C) = E_{ext}(C) + E_{int}(C)$$

leads to curves which are short and stiff while passing through locations of large gradient.

The Mumford-Shah

Euler-Lagrange Equations

Gradient Descent Implementations

The Snakes energy

$$E(C) = -\int_{0}^{1} |\nabla I(C)|^{2} ds + \int_{0}^{1} \left\{ \frac{\alpha}{2} |C_{s}(s)|^{2} + \frac{\beta}{2} |C_{ss}(s)|^{2} \right\} ds$$

is of the canonical form

$$E(C) = \int \mathcal{L}(C, C_s, C_{ss}) ds$$

The corresponding Euler-Lagrange equation is given by:

$$\frac{dE}{dC} = \frac{\partial \mathcal{L}}{\partial C} - \frac{d}{ds} \frac{\partial \mathcal{L}}{\partial C_s} + \frac{d^2}{ds^2} \frac{\partial \mathcal{L}}{\partial C_{ss}} = -\nabla |\nabla I(C)|^2 - \alpha C_{ss} + \beta C_{ssss} = 0.$$

Consequently, the gradient descent equation reads:

$$\frac{\partial C(s,t)}{\partial t} = -\frac{dE(C)}{dC} = \nabla |\nabla I(C)|^2 + \alpha C_{ss} - \beta C_{ssss}$$

The Snakes are among the most influential publications in image processing. To date (Dec '17) they have acquired more than 19000 citations. In 2005, the three authors were awarded an Academy Award for realistic simulations of textiles.

The Snakes are considered the first variational approach to image segmentation.

In comparison to modern segmentation methods, however, they are only of limited practical use:

- Real images typically have many gradient maxima which induce local minima in the cost function E. In practice, the curve must therefore be initialized sufficiently close to the desired solution. Alternatively one can presmooth the input image (to remove spurious local minima). Yet, the smoothing also removes possibly important edge information.
- The evolution of parametric curves is a numerically challenging problem as one needs to avoid self-intersections and instabilities.

Image Segmentation II: Variational Approaches

Dr. Yvain QuÉAU



Variational Image Segmentation

nakes

The Mumford-Shah Model

Euler-Lagrange Equations

Problem with Initialization

Input (square)

and initial curve

Image Segmentation II Variational Approache

Dr. Yvain QuÉAU




Variational Image Seamentation

The Mumford-Shah Model

Euler-Lagrange Equations

Gradient Descent Implementations





smoothed input

final segmentation

Presmoothing makes the edge information "visible" from a larger distance. Yet it removes finer structures like the corners of the square.

(Author: D. Cremers)

 Ballooning: One extends the cost functional by the balloon energy

(L. Cohen, "On active contour models and balloons",

iteratively reducing the smoothness, on each level

& Zisserman, Graduated Non-Convexity, in "Visual

CVGIP 1991), which induces the curve to contract (for

$$E_{balloon}(C) = \gamma \int_{\Omega_{int}(C)} d^2x$$

γ > 0), or to expand (for γ < 0), because the balloon energy simply measures the area of the region Ω_{int} inside the curve *C*.
Coarse-to-fine optimization: One minimizes the Snakes energy starting with a coarse (smoothed image) and

initializing with the previously estimated curve. (See Blake

Reconstruction", 1987).
Global optimization: One reformulates the optimization problem and computes globally optimal solutions (using graph cut methods or convex relaxation methods).

Image Segmentation II: Variational Approaches

Dr. Yvain QuÉAU



Variational Image Segmentation

The Mumford-Shah Model

Euler-Lagrange Equations

Variational Image

The Mumford-Shah

Euler-Lagrange Equations

Gradient Descent

Implementations

Variational Image Segmentation

3 The Mumford-Shah Model

Euler-Lagrange Equations

Variational Image Segmentation

Snakes

The Mumford-Shah

Euler-Lagrange Equations Gradient Descent

Implementations

In 1989, Mumford and Shah proposed to compute a piecewise smooth approximation u of the input image $I:\Omega\subset\mathbb{R}^2\to\mathbb{R}$ by minimizing the functional:

$$E(u,C) = \int_{\Omega} (I(x) - u(x))^2 dx + \lambda \int_{\Omega \setminus C} |\nabla u(x)|^2 dx + \nu |C|,$$

jointly with respect to an approximation $u:\Omega\to\mathbb{R}$ and a one-dimensional discontinuity set $C\subset\Omega$. The three terms have the following meaning:

- The data term assures that u is a faithful approximation of the input I.
- The smoothness term, weighted by λ > 0, assures that u
 is smooth everywhere except for the discontinuity set.
- A further regularizer, weighted by $\nu > 0$, assures that this discontinuity set has minimal length |C|.

For increasing values of the weight λ , the approximation u is forced to be smoother and smoother outside of C. In the limit $\lambda \to \infty$ we obtain a piecewise constant approximation of the image I:

$$E(u,C) = \int_{\Omega} (I(x) - u(x))^2 dx + \nu |C|,$$

where u(x) is constant in each of the regions separated by the boundary C. If we denote these regions by $\{\Omega_1, \ldots, \Omega_n\}$ and the constants by u_i , this can be rewritten as:

$$E(\{u_1,\ldots,u_n\},C) = \sum_{i=1}^n \int_{\Omega_i} (I(x) - u_i)^2 dx + \nu |C|,$$

For the case of two regions, a spatially discrete formulation of this energy is known as the Ising model (Lenz 1920, Ising 1925, Heisenberg 1928). It models the phenomenon of ferromagnetism and is among the most studied models in statistical physics.

Image Segmentation II: Variational Approaches

Dr. Yvain QuÉAU



Variational Image Segmentation


Snakes


The Mumford-Shah

Euler-Lagrange Equations Gradient Descent

Implementations

Discrete Approximation





The length of the curve *C* can be approximated as:

$$|C| \approx \frac{1}{2} \sum_{i \sim j} \left(\frac{u_i - u_j}{2} \right)^2 = \frac{1}{8} \sum_{i \sim j} u_i^2 + u_j^2 - 2u_i u_j = \text{const} - \frac{1}{4} \sum_{i \sim j} u_i u_j,$$

with summation over neighboring pixels i and j. This leads to:

$$E(u) = \sum_{i} (I_i - u_i)^2 - \frac{\nu}{4} \sum_{i \sim i} u_i u_j.$$

E. Ising, "Beitrag zur Theorie des Ferromagnetismus", 1925.

Image Segmentation II: Variational Approaches

Dr. Yvain QuÉAU



Variational Image Segmentation

Snakes


The Mumford-Sha

Euler-Lagrange Equations

Gradient Descent

Implementations

Solution via Graph Cuts



Minimization of the discrete two-region model using graph cuts

Image Segmentation II Variational Approache

Dr. Yvain QuÉAU



Variational Image Segmentation

Snakes

he Mumford-Shah

Euler-Lagrange Equations

Ernst Ising





1925 1995

Ernst Ising * 1900 in Cologne, † 1998 in Peoria, Illinois)

Doctoral thesis with Wilhelm Lenz in Hamburg, emigration to the US in 1947. Image Segmentation II: Variational Approaches

Dr. Yvain QuÉAU



Variational Image Segmentation

Snakes

The Mumford-Shah Model

Euler-Lagrange Equations

The Piecewise Constant Mumford-Shah

For $\lambda \to \infty$ one obtains a special case of the Mumford-Shah functional known as the piecewise constant approximation:

$$E({u_1,\ldots,u_n},C) = \sum_{i=1}^n \int_{\Omega_i} (I(x)-u_i)^2 dx + \nu |C|,$$

This functional is of interest for several reasons:

- It is the spatially continuous version of the discrete spin models (Lenz 1920, Ising 1925, Potts 1956, ...).
 - It is quite powerful, yet mathematically well understood.

In the following, we will therefore discuss several aspects of this functional in more detail:

- Some important mathematical results.
- Euler-Lagrange equations and possible implementations.
 - A statistical interpretation.

Image Segmentation II Variational Approache

Dr. Yvain QuÉAU



Variational Image Seamentation

Snakes

Euler-Lagrange **Equations**

in each region:

Snakes

The Mumford-Shah

Euler-Lagrange Equations Gradient Descent

Implementations

 $\frac{\partial E}{\partial u_i} = 2 \int_{\Omega_i} (I(x) - u_i) dx = 0 \implies u_i = \frac{\int_{\Omega_i} I(x) dx}{\int_{\Omega_i} dx}$

For a given boundary C, the minimizing constants u_i are uniquely determined. They are given by the average brightness

As a result, the cost function is merely a function of the boundary C: $E(C) \equiv \min_u E(C, u)$. In particular, the segmentation has the same mean intensity as the input image.

Existence of minima: There exist minima of the functional E(C). The minimizing boundaries C are closed and differentiable up to discontinuities of the following type:

- Three boundary segments meet at equal angles (120°).
- The boundary meets the domain boundary at a 90° angle.

Minima of E(C) are generally not unique.

Snakes The Mumford-Shah

Model

Euler-Lagrange

Gradient Descent

Implementations

The Mumford-Shah Model

Variational Image Segmentation

4 Euler-Lagrange Equations

Variational Image Segmentation

Snakes

The Mumford-Shah Model

uler-Lagrange

Gradient Descent
Implementations

Unfortunately, the Mumford-Shah functional in its original formulation is not in a canonical form, since the variable of interest (the boundary *C*) appears in the integrand.

There exists an entire research community dedicated to such optimization problems known as shape optimization or shape sensitivity analysis.

In the following we will derive the Euler-Lagrange equation using Green's theorem (following Zhu & Yuille, "Region competition: Unifying snakes, region growing, and Bayes/MDL for multiband image segmentation", PAMI 95).

Assume we are given an energy of the form

$$E(C) = \int_{int(C)} f(x, y) dx dy,$$

where int(C) denotes the region inside a curve C. Let $C: [0,1] \to \mathbb{R}^2$ be a parametric closed curve, with C(s) = (x(s), y(s)).

Snakes

The Mumford-Shah Model

Euler-Lagrange Equations

Gradient Descent
Implementations

Green's Theorem: For a vector field of the form

 $\vec{v}=(a(x,y),\,b(x,y))\in\mathbb{R}^2$ and a closed boundary $\pmb{C}\subset\Omega$ we have:

$$\int_{int(C)} (\nabla \times \vec{v}) d^2 x = \int_{C} \vec{v} ds,$$

where the rotation of v is defined as $\nabla \times \vec{v} \equiv \partial_x b - \partial_y a$. Thus:

$$\int\limits_{int(C)}(b_x-a_y)dxdy=\int\limits_{C}adx+bdy$$

Chosing a vector field \vec{v} such that $f = (b_x - a_y)$, we can rewrite the energy in the canonical form:

$$E(C) = \int_{c} f \, dx dy = \int_{c} a dx + b dy = \int_{0}^{c} (a\dot{x} + b\dot{y}) ds \equiv \int_{0}^{c} \mathcal{L}(x, \dot{x}, y, \dot{y}) ds.$$

where $\dot{x} \equiv \frac{dx(s)}{ds}$ and $\dot{y} \equiv \frac{dy(s)}{ds}$.

Dr. Yvain QuÉAU

Variational Image Segmentation

Snakes

The Mumford-Shah Model

Euler-Lagrange

Gradient Descent
Implementations

The functional

$$E(C) = \int_{int(C)} f(x,y) \, dxdy = \int_{0}^{1} (a\dot{x} + b\dot{y}) ds \equiv \int_{0}^{1} \mathcal{L}(x,\dot{x},y,\dot{y}) ds.$$

is equal to an integral along the curve C and we can compute the functional derivative with respect to C(s) = (x(s), y(s)):

$$\frac{\partial \mathcal{L}}{\partial x} - \frac{d}{ds} \frac{\partial \mathcal{L}}{\partial \dot{x}} = \frac{\partial a}{\partial x} \dot{x} + \frac{\partial b}{\partial x} \dot{y} - \frac{d}{ds} a = \left(\frac{\partial b}{\partial x} - \frac{\partial a}{\partial y}\right) \dot{y} = f \dot{y}$$

$$\frac{\partial \mathcal{L}}{\partial y} - \frac{d}{ds} \frac{\partial \mathcal{L}}{\partial \dot{y}} = \frac{\partial a}{\partial y} \dot{x} + \frac{\partial b}{\partial y} \dot{y} - \frac{d}{ds} b = \left(-\frac{\partial b}{\partial x} + \frac{\partial a}{\partial y}\right) \dot{x} = -f \dot{x}$$

In summary we obtain the simple functional gradient:

$$\frac{dE}{dC} = f(x, y) \begin{pmatrix} \dot{y} \\ -\dot{x} \end{pmatrix} = f(x, y) \vec{n}_C, \text{ where } \vec{n}_C = \text{outer normal}$$

int(C)

Snakes

The Mumford-Shah Model

Euler-Lagrange Equations

Gradient Descent
Implementations

The above calculation shows that functionals of the form $E(C) = \int f(x, y) dxdy$ have the functional derivative:

$$\frac{dE}{dC} = f(x, y)\vec{n}_C.$$

For the piecewise constant Mumford-Shah functional (without boundary length term) and only two regions separated by a curve C we have:

$$E(C) = \int_{int(C)} (I(x) - u_{int})^2 d^2x + \int_{ext(C)} (I(x) - u_{ext})^2 d^2x,$$

so the functional derivative is given by:

$$\frac{dE}{dC} = \left((I(x) - u_{int})^2 - (I(x) - u_{ext})^2 \right) \vec{n}_C,$$

because both regions contribute to the gradient and the outer normal for the outside region is simply $-\vec{n}_C$.

Variational Image Segmentation

The Mumford-Shah Model

Euler-Lagrange Equations

Gradient Descent

Snakes

The Mumford-Shah Model


Euler-Lagrange Equations

Gradient Descent

Implementations

The gradient descent equation is therefore:

$$\frac{\partial C(s,t)}{\partial t} = -\frac{dE(C)}{dC} = \left((I - u_{\text{ext}})^2 - (I - u_{\text{int}})^2 \right) \vec{n}_C.$$



At each boundary point:

Displace the curve

- outwards, if $|I u_{int}| < |I u_{ext}|$
- inwards, if $|I u_{int}| > |I u_{ext}|$

Intuition (Zhu & Yuille, Region Competition, PAMI '96):

If the local brightness I(x) at point x is more similar to the average brightness of the interior then x is assigned to the interior (the curve moves outward) and vice versa.

Snakes

The Mumford-Shah Model

Euler-Lagrange Equations

Gradient Descent

Implementations

For the two-region piecewise constant Mumford-Shah with length regularity we get:

$$E(C) = \int_{int(C)} (I(x) - u_{int})^2 d^2x + \int_{ext(C)} (I(x) - u_{ext})^2 d^2x + \nu |C|$$

and the gradient descent reads:

$$\frac{\partial \textit{C}(\textit{s},\textit{t})}{\partial \textit{t}} = -\frac{\textit{dE}(\textit{C})}{\textit{dC}} = \left((\textit{I} - \textit{u}_{\textit{ext}})^2 - (\textit{I} - \textit{u}_{\textit{int}})^2 - \nu \kappa_{\textit{C}} \right) \vec{\textit{n}}_{\textit{C}},$$

where κ_C denotes the local curvature of the curve C.

This means that in addition to separating bright and dark areas, the evolution aims at suppressing large curvature of the curve. This is what leads to a local minimization of the boundary length $|\mathcal{C}|$.

Snakes

The Mumford-Shah Model

Euler-Lagrange Equations

Gradient Descent

Implementations

Variational Image Segmentation

The Mumford-Shah Model

Euler-Lagrange Equations

Implementations

The paper of Mumford and Shah is focused on aspects of existence and uniqueness of solutions and the study of properties of solutions. It does not propose a numerical implementation for finding minimizers.

There exist a number of alternative methods, for example:

- Ambrosio and Tortorelli, "Approximation of functionals depending on jumps by elliptic functionals via Γ-convergence", Comm. Pure Appl. Math. 90: an approximation of the piecewise smooth model using quadratic functionals.
- Koepfler et al, "A multiscale algorithm for image segmentation by variational method", SIAM JNA 94: an implementation of the piecewise constant model in the spirit of region merging based on the notion of 2-normal segmentations.
- Cremers et al., "Diffusion snakes: Introducing statistical shape knowledge into the Mumford-Shah functional", IJCV 2002: Implementation of the piecewise smooth and piecewise constant models using closed parametric spline curves (hybrid of the Mumford-Shah and the Snakes).

Image Segmentation II: Variational Approaches

Dr. Yvain QuÉAU



Variational Image Segmentation

Snakes

The Mumford-Shah Model

Euler-Lagrange Equations

Gradient Descent

updated 2017-12-06 30/43

Ambrosio and Tortorelli showed in 1990 that the piecewise-smooth Mumford-Shah model

$$E(u,C) = \iint_{\mathbb{R}} \left(I(x) - u(x) \right)^2 dx + \lambda \iint_{\mathbb{R}} |\nabla u(x)|^2 dx + \nu \iint_{\mathbb{R}} |C'(s)| ds$$

can be approximated well by

$$E_{\epsilon}(u, w) = \iint_{\Omega} (I(x) - u(x))^{2} dx + \lambda \iint_{\Omega} w(x)^{2} |\nabla u(x)|^{2} dx$$
$$+ \nu \iint_{\Omega} \epsilon |\nabla w(x)|^{2} + \frac{1}{4\epsilon} [w(x) - 1]^{2} dx$$

in the sense that $E_{\epsilon} \stackrel{\Gamma}{\to} E$ when $\epsilon \to 0$, with $w : \Omega \to \mathbb{R}$ converging towards the indicator function of the the "smooth" part $\Omega \setminus C$.

Alternating optimization then boils down to a sequence of simple linear least-squares problems!

Image Segmentation II Variational Approaches

Dr. Yvain QuÉAU



Snakes

The Mumford-Shah

Euler-Lagrange **Equations**

Gradient Descent

Ambrosio-Tortorelli Approximation

Results with $\lambda = 10$, $\nu = 10^{-7}$ and $\epsilon = 10^{-6}$:







W




Image Segmentation II: Variational Approaches

Dr. Yvain QuÉAU



Variational Image Segmentation

Snakes

The Mumford-Shah Model

Euler-Lagrange Equations

Gradient Descent

mplementations

2-normal Segmentations

Definition (Koepfler et al.'95):

A 2-normal segmentation is a partitioning of the image plane Ω into pairwise disjoint regions $\Omega_1, \ldots, \Omega_n$, such that each segmentation obtained by merging two neighboring regions has a larger or equal energy (in the sense of the piecewise constant Mumford-Shah energy).

The algorithm of Koepfler et al. allows to compute 2-normal segmentations. To this end it iteratively merges neighboring regions until convergence.

Minima of the piecewise constant Mumford-Shah are always 2-normal segmentations.

<u>However:</u> Not all 2-normal segmentations are minimizers of the Mumford-Shah functional.

Two questions arise:

- In which order should one merge neighboring regions?
- How should one select the parameter ν ?

Image Segmentation II: Variational Approaches

Dr. Yvain QuÉAU



Variational Image Segmentation

Snakes

The Mumford-Shah Model

Euler-Lagrange Equations Gradient Descent

nplementations

Proposition: The number n of regions of a 2-normal segmentation is bounded by the following function of the scale parameter ν :

$$n \leq \frac{|\Omega| \operatorname{osc}(I)^4}{c_0 \, \nu^2}.$$

 $|\Omega| \equiv \text{image size}$, and $\operatorname{osc}(I) \equiv \sup(I) - \inf(I)$ is called the oscillation of the brightness function (difference between largest and smallest brightness).

In particular, this implies: The smaller Ω and the larger ν , the stronger the constraint on the number of regions.

Thus, the scale parameter ν defines the spatial scale on which segmentation is performed – on a coarse scale for large ν and on a finer scale for smaller values of ν .

Image Segmentation II: Variational Approaches

Dr. Yvain QuÉAU



Variational Image Segmentation

Snakes

The Mumford-Shah Model

Euler-Lagrange Equations Gradient Descent

nplementations

Properties of 2-normal Segmentations

Moreover, one can compute bounds on the size and shape of segments.

Proposition: The individual segments of a 2-normal segmentation have a a positive minimal size:

$$|\Omega_i| \geq c_1(I, \nu, \Omega) \quad \forall i.$$

In particular, this implies that the Mumford-Shah segmentation process (for $\nu>0$) leads to an elimination of small regions. In historical approaches this was introduced through a heuristic post-processing step.

Proposition: For every individual segment Ω_i the length of its boundary $\partial \Omega_i$ is bounded by a mutiple of its area $|\Omega_i|$:

$$|\partial\Omega_i|\leq c_2(I,\Omega)|\Omega_i|\quad\forall i.$$

This implies that minimization of the Mumford-Shah functional (for $\nu > 0$) also leads to an elimination of elongated regions.

Image Segmentation II: Variational Approaches

Dr. Yvain QuÉAU



Variational Image Segmentation

Snakes

The Mumford-Shah Model

Euler-Lagrange Equations Gradient Descent

plementations

updated 2017-12-06 35/43

Snakes

Model

Euler-Lagrange Equations

The Mumford-Shah

Gradient Descent

The implementation of Koepfler et al. allows to compute a 2-normal segmentation by the following region merging process:

- Initialize: every pixel is its own region.
- 2 For all neighboring pairs of regions, compute the change ΔE in energy obtained by merging the two regions. Obviously it is of the form:

$$\Delta E = \Delta E_{region} + \nu \Delta E_{length}$$

- 3 For all pairs of adjacent regions determine the value $\hat{\nu}$, for which there is an energy decrease ($\Delta E < 0$). This value always exists because $\Delta E_{length} < 0$ and $\Delta E_{region} \ge 0$.
- 4 In each step, merge the region pair with the smallest value of $\hat{\nu}$.
- 5 Repeat steps (2.) (4.) until the desired number of regions or a sufficiently large value of ν is reached.

Multiscale Implementation

Input

Segmentations

Segmentations for $\nu = 2022$ (above) and $\nu = 6173$ (below)

Koepfler et al. '95

http://www.math-info.univ-paris5.fr/~qk/papers/heidelberg95.pdf

Image Segmentation II Variational Approache

Dr. Yvain QuÉAU



Variational Image Segmentation

Snakes

The Mumford-Shah Model

Euler-Lagrange Equations Gradient Descent

nplementations

Multiscale Implementation

Image Segmentation II: Variational Approaches

Dr. Yvain QuÉAU



Variational Image Segmentation

Snakes

The Mumford-Shah Model

Euler-Lagrange Equations

Gradient Descent

nplementations



Original and segmentation with $\nu = 1024$ and $\nu = 4096$

Koepfler, Morel, Solimini, "Segmentation by minimizing functionals and the merging methods", GRETSI '91

$$E(u,C) = \int_{\Omega} (I(x) - u(x))^2 dx + \lambda \int_{\Omega \setminus C} |\nabla u(x)|^2 dx + \nu \int_{0}^{1} |C'(s)|^2 ds$$

by alternating two gradient descent evolutions:

$$\frac{\partial C(s,t)}{\partial t} = -\frac{\partial E}{\partial C} = \left((I - u)^2 + \lambda |\nabla u|^2 \right)_-^+ \vec{n} + 2\nu C'', \quad \vec{n} = \text{normal}$$

$$\frac{\partial u(x,t)}{\partial t} = -\frac{\partial E}{\partial u} = \lambda \nabla (w_c \nabla u) + (I - u), \quad w_c(x) = \begin{cases} 0, & x \in C \\ 1, & \text{else} \end{cases}$$

Here f_{-}^{+} denotes the difference of integrand f on either side of the boundary – see slide 25. This gives a curve evolution in alternation with an inhomogeneous diffusion process (constant diffusion inside regions, no diffusion across boundary). Thus the Diffusion Snakes simultaneously perform denoising (in each of the separated regions) and boundary estimation.

Image Segmentation II Variational Approaches

Dr. Yvain QuÉAU



Seamentation

Snakes

The Mumford-Shah Euler-Lagrange

Equations


Gradient Descent

Diffusion Snakes


The evolution of the curve C is implemented by evolving a finite number of control points $p_1, \ldots, p_n \in \mathbb{R}^2$:

$$C(s,t) = \sum_{i=1}^n p_i(t) B_i(s).$$

Here $B_i(s)$ are spline basis functions:



spline basis functions



Spline & control points

Image Segmentation II: Variational Approaches

Dr. Yvain QuÉAU



Variational Image Segmentation

Snakes

The Mumford-Shah Model

Euler-Lagrange Equations

Gradient Descent

nnlementations

Diffusion Snakes: Curve Evolution

Image Segmentation II: Variational Approaches

Dr. Yvain QuÉAU



Variational Image Segmentation

Snakes

The Mumford-Shah Model

Euler-Lagrange Equations

Gradient Descent

mplementations

Cremers et al., "Diffusion snakes: Introducing statistical shape knowledge into the Mumford-Shah functional", IJCV 2002

Diffusion Snakes



 ν large

 ν small

Cremers et al., "Diffusion snakes: Introducing statistical shape knowledge into the Mumford-Shah functional", IJCV 2002

Image Segmentation II: Variational Approaches

Dr. Yvain QuÉAU



Variational Image Segmentation

Snakes

The Mumford-Shah Model

Euler-Lagrange Equations Gradient Descent

nplementations

Segmentation with Statistical Shape Priors

Image Segmentation II: Variational Approaches

Dr. Yvain QuÉAU



Variational Image Segmentation

Snakes

The Mumford-Shah Model

Euler-Lagrange Equations Gradient Descent

mplementations

 $E(C) = E_{diffsnake}(C) + E_{prior}(C) \rightarrow \min$

Cremers et al., "Diffusion snakes: Introducing statistical shape knowledge into the Mumford-Shah functional". IJCV 2002