

Chapter 1

Convex Analysis

Convex Optimization for Machine Learning & Computer Vision
WS 2018/19

Tao Wu
Yuesong Shen
Zhenzhang Ye

Computer Vision Group
Department of Informatics
TU Munich

Convex Analysis

Tao Wu
Yuesong Shen
Zhenzhang Ye



Convex Set

Convex Function

Existence of Minimizer

Subdifferential



Convex Set

Notations

- \mathbb{E} is a *Euclidean space* (i.e., finite dimensional inner product space), equipped with

① Inner product $\langle \cdot, \cdot \rangle$, e.g., $\langle u, v \rangle = u^\top v$ if $\mathbb{E} = \mathbb{R}^n$;

② Norm $\|\cdot\| = \sqrt{\langle \cdot, \cdot \rangle}$ satisfying polarization identity:

$$2\|u\|^2 + 2\|v\|^2 = \|u + v\|^2 + \|u - v\|^2.$$

- C is a closed, convex subset of \mathbb{E} .
- J is a convex *objective* function.



Convex Set

Convex Function

Existence of Minimizer

Subdifferential



Notations

- \mathbb{E} is a *Euclidean space* (i.e., finite dimensional inner product space), equipped with

① Inner product $\langle \cdot, \cdot \rangle$, e.g., $\langle u, v \rangle = u^\top v$ if $\mathbb{E} = \mathbb{R}^n$;

② Norm $\|\cdot\| = \sqrt{\langle \cdot, \cdot \rangle}$ satisfying polarization identity:

$$2\|u\|^2 + 2\|v\|^2 = \|u + v\|^2 + \|u - v\|^2.$$

- C is a closed, convex subset of \mathbb{E} .
- J is a convex *objective* function.

Convex Set

Convex Function

Existence of Minimizer

Subdifferential

Convex optimization

$$\text{minimize } J(u) \quad \text{over } u \in C.$$

First questions:

- What is a convex set?
- What is a convex function?

Convex set

Definition

A set C is said to be **convex** if

$$\alpha u + (1 - \alpha)v \in C, \quad \forall u, v \in C, \quad \forall \alpha \in [0, 1].$$




Convex Set

Convex Function


Existence of Minimizer

Subdifferential

convex



non-convex



Definition

- A set $C \subset \mathbb{E}$ is **open** if $\forall u \in C, \exists \epsilon > 0$ s.t. $B_\epsilon(u) \subset C$, where $B_\epsilon(u) := \{v \in \mathbb{E} : \|v - u\| < \epsilon\}$.
- A set $C \subset \mathbb{E}$ is **closed** if its complement $\mathbb{E} \setminus C$ is open.
- The **closure** of a set $C \subset \mathbb{E}$ is

$$\text{cl } C = \{u \in \mathbb{E} : \exists \{u^k\} \subset C \text{ s.t. } \lim_{k \rightarrow \infty} u^k = u\}.$$

- The **interior** of a set $C \subset \mathbb{E}$ is

$$\text{int } C = \{u \in C : \exists \epsilon > 0 \text{ s.t. } B_\epsilon(u) \subset C\}.$$





Definition

- A set $C \subset \mathbb{E}$ is **open** if $\forall u \in C, \exists \epsilon > 0$ s.t. $B_\epsilon(u) \subset C$, where $B_\epsilon(u) := \{v \in \mathbb{E} : \|v - u\| < \epsilon\}$.
- A set $C \subset \mathbb{E}$ is **closed** if its complement $\mathbb{E} \setminus C$ is open.
- The **closure** of a set $C \subset \mathbb{E}$ is

$$\text{cl } C = \{u \in \mathbb{E} : \exists \{u^k\} \subset C \text{ s.t. } \lim_{k \rightarrow \infty} u^k = u\}.$$

- The **interior** of a set $C \subset \mathbb{E}$ is

$$\text{int } C = \{u \in C : \exists \epsilon > 0 \text{ s.t. } B_\epsilon(u) \subset C\}.$$

- The **relative interior** of a set $C \subset \mathbb{E}$ is

$$\begin{aligned} \text{rint } C &:= \{u \in C : \exists \epsilon > 0 \text{ s.t. } B_\epsilon(u) \cap \text{aff } C \subset C\} \\ &= \{u \in C : \forall v \in C, \exists \alpha > 1 \text{ s.t. } v + \alpha(u - v) \in C\} \end{aligned}$$

if C is convex. Here $\text{aff } C$ stands for the **affine hull** of C .



The following operations preserve the convexity:

- Intersection: $C_1 \cap C_2$.
- Summation: $C_1 + C_2 := \{u^1 + u^2 : u^1 \in C_1, u^2 \in C_2\}$.
- Closure: $\text{cl } C$.
- (Relative) interior: $\text{int } C$, $\text{rint } C$.

– In general, the union of convex sets is not convex.



The following operations preserve the convexity:

- Intersection: $C_1 \cap C_2$.
- Summation: $C_1 + C_2 := \{u^1 + u^2 : u^1 \in C_1, u^2 \in C_2\}$.
- Closure: $\text{cl } C$.
- (Relative) interior: $\text{int } C$, $\text{rint } C$.

– In general, the union of convex sets is not convex.

Convex cone

C is a **cone** if $C = \alpha C$ for any $\alpha > 0$.

C is a **convex cone** if C is a cone and is convex as well.

Separation of convex sets

Theorem (separation of convex sets)

Let C_1, C_2 be nonempty convex subsets in \mathbb{E} .

- ① Assume C_1 is closed and $C_2 = \{w\} \subset \mathbb{E} \setminus C_1$. Then $\exists v \in \mathbb{E}, v \neq 0, \alpha \in \mathbb{R}$ s.t.

$$\langle v, w \rangle > \alpha \geq \langle v, u \rangle, \quad \forall u \in C_1.$$

- ② Assume C_1 is open and $C_2 = \{w\} \subset \mathbb{E} \setminus C_1$. Then $\exists v \in \mathbb{E}, v \neq 0, \alpha \in \mathbb{R}$ s.t.

$$\langle v, w \rangle \geq \alpha \geq \langle v, u \rangle, \quad \forall u \in C_1.$$

- ③ Assume $C_1 \cap C_2 = \emptyset$ and C_1 is open. Then $\exists v \in \mathbb{E}, v \neq 0, \alpha \in \mathbb{R}$ s.t.

$$\langle v, u^1 \rangle \geq \alpha \geq \langle v, u^2 \rangle, \quad \forall u^1 \in C_1, u^2 \in C_2.$$

- ④ Assume $\emptyset \neq \text{int } C_1 \subset \mathbb{E} \setminus C_2$. Then $\exists v \in \mathbb{E}, v \neq 0, \alpha \in \mathbb{R}$ s.t.

$$\langle v, u^1 \rangle \geq \alpha \geq \langle v, u^2 \rangle, \quad \forall u^1 \in C_1, u^2 \in C_2.$$



Convex Set

Convex Function

Existence of Minimizer

Subdifferential



Convex Function



- An **extended real-valued function** J maps from \mathbb{E} to $\bar{\mathbb{R}} := \mathbb{R} \cup \{\infty\}$.
- The **domain** of $J : \mathbb{E} \rightarrow \bar{\mathbb{R}}$ is

$$\text{dom } J = \{u \in \mathbb{E} : J(u) < \infty\}.$$

- The function $J : \mathbb{E} \rightarrow \bar{\mathbb{R}}$ is **proper** if $\text{dom } J \neq \emptyset$.

Definition

We say $J : \mathbb{E} \rightarrow \bar{\mathbb{R}}$ is a **convex function** if

- 1 $\text{dom } J$ is a convex set.
- 2 For all $u, v \in \text{dom } J$ and $\alpha \in [0, 1]$ it holds that

$$J(\alpha u + (1 - \alpha)v) \leq \alpha J(u) + (1 - \alpha)J(v).$$

We say J is **strictly convex** if the above inequality is strict for all $\alpha \in (0, 1)$ and $u \neq v$.

Examples

- $J_{data}(u) = \|u - z\|_q^q$, where $q \geq 1$ and $\|\cdot\|_q$ is ℓ^q -norm.
- $J_{regu}(u) = \|Ku\|_{q'}^{q'}$, where K is linear transform and $q' \geq 1$.
- $J(u) = J_{data}(u) + \alpha J_{regu}(u)$.



Examples

- $J_{data}(u) = \|u - z\|_q^q$, where $q \geq 1$ and $\|\cdot\|_q$ is ℓ^q -norm.
- $J_{regu}(u) = \|Ku\|_{q'}^{q'}$, where K is linear transform and $q' \geq 1$.
- $J(u) = J_{data}(u) + \alpha J_{regu}(u)$.
- Negative entropy: $J_\epsilon(u) = \epsilon(u \log(u) + (1 - u) \log(1 - u))$.
- Soft plus: $J_\epsilon^*(v) = \epsilon \log(1 + \exp(v/\epsilon)) \xrightarrow{\epsilon \rightarrow 0^+} \max(v, 0)$.



Examples

- $J_{data}(u) = \|u - z\|_q^q$, where $q \geq 1$ and $\|\cdot\|_q$ is ℓ^q -norm.
- $J_{regu}(u) = \|Ku\|_{q'}^{q'}$, where K is linear transform and $q' \geq 1$.
- $J(u) = J_{data}(u) + \alpha J_{regu}(u)$.
- Negative entropy: $J_\epsilon(u) = \epsilon(u \log(u) + (1 - u) \log(1 - u))$.
- Soft plus: $J_\epsilon^*(v) = \epsilon \log(1 + \exp(v/\epsilon)) \xrightarrow{\epsilon \rightarrow 0^+} \max(v, 0)$.
- **Indicator function** ($C \subset \mathbb{E}$ is closed and convex):

$$\delta_C(u) = \begin{cases} 0 & \text{if } u \in C, \\ \infty & \text{otherwise.} \end{cases}$$

- Formulate *constrained optimization* with indicator function:
 $\min J(u)$ over $u \in C$. \Leftrightarrow $\min J(u) + \delta_C(u)$ over $u \in \mathbb{E}$.





(As exercises)

- Any norm (over a normed vector space) is a convex function.
- J is a convex function and A is an affine transform $\Rightarrow J(A \cdot)$ is convex function.
- (Jensen's inequality) $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ is convex iff

$$J\left(\sum_{i=1}^n \alpha_i u^i\right) \leq \sum_{i=1}^n \alpha_i J(u^i),$$


whenever $\{u^i\}_{i=1}^n \subset \mathbb{E}$, $\{\alpha_i\}_{i=1}^n \subset [0, 1]$, $\sum_{i=1}^n \alpha_i = 1$.

Epigraph

Definition

The **epigraph** of a proper function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ is

$$\text{epi } J = \{(u, \alpha) \in \mathbb{E} \times \mathbb{R} : J(u) \leq \alpha\}.$$



Theorem

A proper function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ is convex (resp. strictly convex) iff $\text{epi } J$ is a convex (resp. strictly convex) set.

Proof: as exercise.



Definition

Assume $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ with $\text{rint dom } J \neq \emptyset$. We say J is **locally Lipschitz** at $u \in \text{rint dom } J$ with modulus $L_u > 0$ if there exists $\epsilon > 0$ s.t.

$$|J(u^1) - J(u^2)| \leq L_u \|u^1 - u^2\| \quad \forall u^1, u^2 \in B_\epsilon(u) \cap \text{rint dom } J.$$



Convex Set

Convex Function

Existence of Minimizer

Subdifferential



Definition

Assume $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ with $\text{rint dom } J \neq \emptyset$. We say J is **locally Lipschitz** at $u \in \text{rint dom } J$ with modulus $L_u > 0$ if there exists $\epsilon > 0$ s.t.

$$|J(u^1) - J(u^2)| \leq L_u \|u^1 - u^2\| \quad \forall u^1, u^2 \in B_\epsilon(u) \cap \text{rint dom } J.$$

Theorem

A proper convex function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ is locally Lipschitz at any $u \in \text{rint dom } J$.

Proof: found in script.

Convex Set

Convex Function

Existence of Minimizer

Subdifferential



Existence of Minimizer

Global vs. Local minimizer

Recall the optimization of $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$:

$$\text{minimize } J(u) \quad \text{over } u \in \mathbb{E}.$$



Definition

- 1 $u^* \in \mathbb{E}$ is a **global minimizer** if $J(u^*) \leq J(u)$ for all $u \in \mathbb{E}$.
- 2 u^* is a **local minimizer** if $\exists \epsilon > 0$ s.t. $J(u^*) \leq J(u)$ for all $u \in B_\epsilon(u^*)$.
- 3 In the above definitions, a global/local minimizer is **strict** if $J(u^*) \leq J(u)$ is replaced by $J(u^*) < J(u)$.

Global vs. Local minimizer

Recall the optimization of $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$:

$$\text{minimize } J(u) \quad \text{over } u \in \mathbb{E}.$$



Definition

- 1 $u^* \in \mathbb{E}$ is a **global minimizer** if $J(u^*) \leq J(u)$ for all $u \in \mathbb{E}$.
- 2 u^* is a **local minimizer** if $\exists \epsilon > 0$ s.t. $J(u^*) \leq J(u)$ for all $u \in B_\epsilon(u^*)$.
- 3 In the above definitions, a global/local minimizer is **strict** if $J(u^*) \leq J(u)$ is replaced by $J(u^*) < J(u)$.

Theorem

For any proper convex function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$, if $u^* \in \text{dom } J$ is a local minimizer of J , then it is also a global minimizer.

Proof: on board.

Does a minimizer always exist?


- Consider

$$\text{minimize } J(u) \quad \text{over } u \in \mathbb{E},$$


where $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ is a proper, convex function.

- Some counterexamples for $J : \mathbb{R} \rightarrow \overline{\mathbb{R}}$:


u



$\exp u$



$u^2 + \delta\{u > 0\}$



- Next we formalize our observations and derive sufficient conditions for existence.



Definition

- 1 J is **bounded from below** if $J(\cdot) \geq C$ for some $C \in \mathbb{R}$.
- 2 J is **coercive** if $J(u) \rightarrow \infty$ whenever $\|u\| \rightarrow \infty$.
- 3 J is **lower semi-continuous** (lsc) at u^* if

$$J(u^*) \leq \liminf_{k \rightarrow \infty} J(u^k), \text{ whenever } u^k \rightarrow u^*.$$

Theorem

Any proper function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$, which is bounded from below, coercive, and lsc (everywhere), has a (global) minimizer.

Proof: on board.



Convex Set

Convex Function

Existence of Minimizer

Subdifferential



- Recall that a function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ is strictly convex if

$$J(\alpha u + (1 - \alpha)v) < \alpha J(u) + (1 - \alpha)J(v),$$

for all $u, v \in \text{dom } J$, $u \neq v$, $\alpha \in (0, 1)$.


Theorem

The minimizer of a strictly convex function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ is unique.

Proof: on board.



Subdifferential




Definition

$J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ is called (Fréchet) **differentiable** at $u \in \text{int dom } J$ and $\nabla J(u) \in \mathbb{E}$ is the (Fréchet) **differential** of J at u if

$$\lim_{h \rightarrow 0} \frac{|J(u+h) - J(u) - \langle \nabla J(u), h \rangle|}{\|h\|} = 0.$$

J is **continuously differentiable** at $u \in \text{int dom } J$ if $\nabla J(\cdot)$ is continuous on $(\text{dom } J) \cap B_\epsilon(u)$ for some $\epsilon > 0$.





Definition

$J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ is called (Fréchet) **differentiable** at $u \in \text{int dom } J$ and $\nabla J(u) \in \mathbb{E}$ is the (Fréchet) **differential** of J at u if

$$\lim_{h \rightarrow 0} \frac{|J(u+h) - J(u) - \langle \nabla J(u), h \rangle|}{\|h\|} = 0.$$

J is **continuously differentiable** at $u \in \text{int dom } J$ if $\nabla J(\cdot)$ is continuous on $(\text{dom } J) \cap B_\epsilon(u)$ for some $\epsilon > 0$.

Remark

If \mathbb{E} is a topological vector space, $\nabla J(u)$ is treated as a *dual* object in \mathbb{E}^* , and $\langle \nabla J(u), h \rangle_{\mathbb{E}^*, \mathbb{E}}$ as *duality pairing*.




Subdifferential

Definition

The **subdifferential** of a convex function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ at $u \in \text{dom } J$ is defined by

$$\partial J(u) = \{p \in \mathbb{E} : J(v) \geq J(u) + \langle p, v - u \rangle \forall v \in \mathbb{E}\}.$$




Subdifferential

Definition

The **subdifferential** of a convex function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ at $u \in \text{dom } J$ is defined by

$$\partial J(u) = \{p \in \mathbb{E} : J(v) \geq J(u) + \langle p, v - u \rangle \quad \forall v \in \mathbb{E}\}.$$



Geometric interpretation



$p \in \partial J(u)$ iff $(p, -1)$ is a normal vector for the supporting hyperplane of $\text{epi } J$ at $(u, J(u))$, i.e.,

$$\left\langle \begin{bmatrix} p \\ -1 \end{bmatrix}, \begin{bmatrix} u \\ J(u) \end{bmatrix} \right\rangle \geq \left\langle \begin{bmatrix} p \\ -1 \end{bmatrix}, \begin{bmatrix} v \\ \alpha \end{bmatrix} \right\rangle, \quad \forall (v, \alpha) \in \text{epi } J.$$





Subdifferential: Examples

① $J(u) = |u|$.



Subdifferential: Examples

① $J(u) = |u|$.



② Given a closed, convex subset $C \subset \mathbb{E}$ and $u \in C$,



$$\partial\delta_C(u) = \{p \in \mathbb{E} : \langle p, v - u \rangle \leq 0 \forall v \in C\} =: N_C(u),$$

known as the *normal cone* of C at u .



Subdifferential: Examples

① $J(u) = |u|$.



② Given a closed, convex subset $C \subset \mathbb{E}$ and $u \in C$,

$$\partial \delta_C(u) = \{p \in \mathbb{E} : \langle p, v - u \rangle \leq 0 \forall v \in C\} =: N_C(u),$$

known as the *normal cone* of C at u .

③ $J(u) = \|u\| \Rightarrow \partial J(0) = \{p : \|p\|_* \leq 1\}$. $\|\cdot\|_*$ is the dual norm of $\|\cdot\|$, i.e., $\|p\|_* = \sup\{\langle p, u \rangle : \|u\| \leq 1\}$.



Theorem (chain rule under linear transform)

Let $\tilde{J}(\cdot) = J(K\cdot)$ with some convex function J and linear transform K . Then

$$\partial\tilde{J}(u) = K^\top \partial J(Ku)$$

whenever $Ku \in \text{rint dom } J$.

Example: $J(u) = \|Ku\| \Rightarrow \partial J(u) = K^\top \partial \|\cdot\|(Ku)$.



Convex Set

Convex Function

Existence of Minimizer

Subdifferential



Theorem (chain rule under linear transform)

Let $\tilde{J}(\cdot) = J(K\cdot)$ with some convex function J and linear transform K . Then

$$\partial\tilde{J}(u) = K^\top \partial J(Ku)$$

whenever $Ku \in \text{rint dom } J$.

Example: $J(u) = \|Ku\| \Rightarrow \partial J(u) = K^\top \partial \|\cdot\|(Ku)$.

Theorem (summation rule)

If $\tilde{J}(\cdot) = J_1(\cdot) + J_2(\cdot)$ for some convex functions J_1 and J_2 , then

$$\partial\tilde{J}(u) = \partial J_1(u) + \partial J_2(u)$$

for any $u \in \text{rint dom } J_1 \cap \text{rint dom } J_2$.

Properties of subdifferential map

Theorem

Let $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ be a convex function. Then for any $u \in \text{int dom } J$, $\partial J(u)$ is a nonempty, compact, and convex subset.

Proof: on board.

Convex Analysis

Tao Wu
Yuesong Shen
Zhenzhang Ye



Convex Set

Convex Function

Existence of Minimizer

Subdifferential

Properties of subdifferential map

Theorem

Let $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ be a convex function. Then for any $u \in \text{int dom } J$, $\partial J(u)$ is a nonempty, compact, and convex subset.

Proof: on board.

Theorem

Let $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ be a convex function. Then ∂J is a **monotone operator**, i.e. $\forall u^1, u^2 \in \text{dom } J, p^1 \in \partial J(u^1), p^2 \in \partial J(u^2)$:

$$\langle p^1 - p^2, u^1 - u^2 \rangle \geq 0.$$

Proof: on board.



Properties of subdifferential map

Theorem

Let $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ be a convex function. Then for any $u \in \text{int dom } J$, $\partial J(u)$ is a nonempty, compact, and convex subset.

Proof: on board.

Theorem

Let $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ be a convex function. Then ∂J is a **monotone operator**, i.e. $\forall u^1, u^2 \in \text{dom } J, p^1 \in \partial J(u^1), p^2 \in \partial J(u^2)$:

$$\langle p^1 - p^2, u^1 - u^2 \rangle \geq 0.$$

Proof: on board.

Theorem

Let $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ be a proper, convex, lsc function. Then the set-valued map ∂J is **closed**, i.e. $p^* \in \partial J(u^*)$ whenever

$$\exists (u^k, p^k) \rightarrow (u^*, p^*) \in (\text{dom } J) \times \mathbb{E} \text{ s.t. } p^k \in \partial J(u^k) \quad \forall k.$$

Proof: on board.



Optimality condition

Theorem

Given any proper convex function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$, the sufficient and necessary condition for u^* being a (global) minimizer for J is

$$0 \in \partial J(u^*).$$

Proof: on board.



Optimality condition

Theorem

Given any proper convex function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$, the sufficient and necessary condition for u^* being a (global) minimizer for J is


$$0 \in \partial J(u^*).$$

Proof: on board.

Constrained optimization as special case

If u^* minimizes $\tilde{J} = J + \delta_C$ with convex function $J : \mathbb{E} \rightarrow \mathbb{R}$ and closed convex subset $C \subset \mathbb{E}$, then $0 \in \partial \tilde{J}(u^*) \Leftrightarrow$

$$0 \in \partial J(u^*) + N_C(u^*).$$




Optimality condition

Constrained optimization as special case

If u^* minimizes $\tilde{J} = J + \delta_C$ with convex function $J : \mathbb{E} \rightarrow \mathbb{R}$ and closed convex subset $C \subset \mathbb{E}$, then $0 \in \partial \tilde{J}(u^*) \Leftrightarrow$

$$0 \in \partial J(u^*) + N_C(u^*).$$



Remark

The optimality condition $0 \in \partial J(u^*) + N_C(u^*)$ is *geometric*. More explicit characterization relies on the *algebraic* representation of $N_C(u^*)$ (e.g., the **Karush-Kuhn-Tucker (KKT) conditions**) typically under certain *constraint qualifications*.



Optimality condition

Constrained optimization as special case

If u^* minimizes $\tilde{J} = J + \delta_C$ with convex function $J : \mathbb{E} \rightarrow \mathbb{R}$ and closed convex subset $C \subset \mathbb{E}$, then $0 \in \partial \tilde{J}(u^*) \Leftrightarrow$

$$0 \in \partial J(u^*) + N_C(u^*).$$

Remark

The optimality condition $0 \in \partial J(u^*) + N_C(u^*)$ is *geometric*. More explicit characterization relies on the *algebraic* representation of $N_C(u^*)$ (e.g., the **Karush-Kuhn-Tucker (KKT) conditions**) typically under certain *constraint qualifications*.

Example: Linear-inequality constraints

Let $C = \{u \in \mathbb{R}^n : Au \leq b\}$ where $b \in \mathbb{R}^m$, $A \in \mathbb{R}^{m \times n}$ has linearly independent rows. Then

$$N_C(u) = \{A^\top \lambda : \lambda \geq 0, \lambda_i = 0 \text{ if } (Au - b)_i < 0\}.$$

