

## Weekly Exercises 11

Room: 02.09.023

Wednesday, 05.02.2020, 12:15 - 14:00

Submission deadline: Monday, 03.02.2020, 16:15, Room 02.09.023

### 1 Convergence

(6+6 Points)

**Exercise 1** (6 Points). Denote  $\Pi_C(x)$  as the projection of point  $x$  onto a nonempty closed convex set  $C$ . Show following properties:

- $\Pi_C$  is a monotone operator.
- $T$  is firmly nonexpansive, if and only if  $\|Tx - Ty\|^2 \leq \langle x - y, Tx - Ty \rangle$ ,  $\forall x, y$ .
- $\Pi_C$  is firmly nonexpansive.

Hint: you might use that  $\langle y - \Pi_C(x), x - \Pi_C(x) \rangle \leq 0, \forall y \in C$

**Exercise 2** (6 Points). Prove the Krasnoselskii-Mann Theorem: Let  $C$  be a nonempty, closed, convex subset of  $\mathbb{E}$ , and  $u^{k+1} = (1 - \tau^k)u^k + \tau^k\Psi(u^k)$  for  $k = 0, 1, 2, \dots$  where  $\{\tau^k\} \subset [0, 1]$  s.t.

$$\sum_{k=0}^{\infty} \tau^k (1 - \tau^k) = \infty,$$

and  $\Psi : C \rightarrow C$  satisfies:

1.  $\Psi$  is nonexpansive.
2.  $\Psi$  has at least one fixed point.

Then  $\{u^k\}$  converges to a fixed point of  $\Psi$ .

### TV- $\ell_1$ denoising (Due: 10.02)

This time, we have an image with salt-and-pepper noise. Therefore, we use  $\ell_1$  norm for the data term.

$$\operatorname{argmin}_{u \in \mathbb{R}^N} \|u - f\|_1 + \rho \|Ku\|_1$$

where  $f \in \mathbb{R}^N$  is the input image with  $N$  pixels,  $\rho$  is a scalar weighting the smooth regularizer and  $K$  is the gradient operator.

Your task is to apply PDHG to solve above problem.

Hint: for choosing the step size, consider the inequality:

$$\|A\| \leq \sqrt{\max_{1 \leq j \leq n} \sum_{i=1}^n |a_{ij}| \max_{1 \leq i \leq m} \sum_{j=1}^m |a_{ij}|},$$

where  $A \in \mathbb{R}^{m \times n}$ .