

Weekly Exercises 8

Room: 02.09.023

Wednesday, 15.01.2020, 12:15-14:00

Submission deadline: Monday, 13.01.2020, 16:15, Room 02.09.023

Exact Line Search

(14+6 Points)

Exercise 1 (6 Points). Let $Q \in \mathbb{R}^{n \times n}$ be a positive definite symmetric matrix. Prove the following inequality for any vector $x \in \mathbb{R}^n$

$$\frac{(x^\top x)^2}{(x^\top Qx)(x^\top Q^{-1}x)} \geq \frac{4\lambda_n\lambda_1}{(\lambda_n + \lambda_1)^2},$$

where λ_n and λ_1 are, respectively, the largest and smallest eigenvalues of Q .

Exercise 2 (6 Points). Let $Q \in \mathbb{R}^{n \times n}$ be symmetric positive definite, and $b \in \mathbb{R}^n$. As in the previous exercise, denote the eigenvalues of Q as $0 < \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n$. Consider the quadratic function $f : \mathbb{R}^n \rightarrow \mathbb{R}$, $x \mapsto \frac{1}{2}x^\top Qx - b^\top x$ and show gradient descent with exact line search has the following convergence property:

$$\|x^{k+1} - x^*\|_Q^2 \leq \left(\frac{\lambda_n - \lambda_1}{\lambda_n + \lambda_1} \right)^2 \|x^k - x^*\|_Q^2,$$

where $x^* \in \mathbb{R}^n$ denotes the global minimizer of f .

Hint: use the inequality from exercise 1.

Image Denoising (Due Date: 13.01)

(10 Points)

It will be published before 19.12.2019.